

Microbial Hazards in Irrigation Water: Standards, Norms, and Testing to Manage Use of Water in Fresh Produce Primary Production

Mieke Uyttendaele, Lee-Ann Jaykus, Philip Amoah, Alessandro Chiodini, David Cunliffe, Liesbeth Jacxsens, Kevin Holvoet, Lise Korsten, Mathew Lau, Peter McClure, Gertjan Medema, Imca Sampers, and Pratima Rao Jasti

Abstract: Accessibility to abundant sources of high-quality water is integral to the production of safe and wholesome fresh produce. However, access to safe water is becoming increasingly difficult in many parts of the world, and this can lead to the production of fresh produce contaminated with pathogenic microorganisms, resulting in increased risk of human disease. Water, an important raw material in the fresh produce chain, is used in considerable amounts in many operations, including irrigation and application of pesticides and fertilizers, but also as a transport medium and for cooling and washing in postharvest practices. In several reported outbreaks related to uncooked fruit and vegetable products, water has been identified as a likely source of the outbreak. The present study, initiated by the ILSI Europe Emerging Microbiological Issues Task Force in collaboration with 8 other ILSI branches and support of WHO/FAO, was undertaken to review the status of, and provide suggestions for, consideration by different stakeholders on water and sanitation and its impact on food safety and public health. A limited number of guidelines and regulations on water quality for agricultural production are available, and many of them are still heavily based on microbial standards and (debated) parameters such as fecal coliforms. Data gaps have been identified with regard to baseline studies of microbial pathogens in water sources in many regions, the need for agreement on methods and microbial parameters to be used in assessing water quality, the fate of pathogens in water, and their transfer and persistence on irrigated/processed produce.

Keywords: water, irrigation, outbreaks, fresh produce, good practices, testing

Introduction

Fresh produce (fresh fruits and vegetables) consumption has been increasing worldwide for several decades (Leon and others 2009; Betts 2014). The reasons are many, but key is the “healthy eating” advice currently being promoted, where the “5 portions per day” message is being widely advocated. Indeed even 7 portions a day has been recently mentioned. The expansion of the fresh produce market over recent years has resulted in a wide variety of fruit and fresh produce being available throughout the year. There is no

doubt that the consumption of increasing amounts of fresh fruits and vegetables is beneficial to the health of the consumer (Dauchet and others 2005).

Microbiologically, however, there can be some challenges in the production of safe fresh produce (Betts 2014), which has been accompanied by a rise in the number of produce-associated foodborne disease outbreaks. In the United States between 1998 and 2007, fresh produce was involved in 684 outbreaks, resulting in 26735 cases of illness. Proportionally, this equates to 14.8% of outbreaks and 22.8% of outbreak-related cases of all foodborne illnesses in the United States. Salads, vegetables, and fruits were linked to 345, 228 cases and 111 outbreaks, respectively (DeWaal and others 2009; Olaimat and Holley 2012). There has also been an increasing association of food-borne outbreaks with vegetables, juices, and other products in the European Union (EU). These products represented 8.7% of reported outbreaks for which a food vehicle was identified in 2010, versus 2.1% in 2009 (EFSA 2012). Based upon EU Zoonoses Monitoring data from 2007 to 2011, Foods of Non-Animal Origin (FoNAO) were associated with 10% of outbreaks, 26% of cases, 35% of hospitalizations, and 46% of deaths (EFSA Panel on Biological Hazards (BIOHAZ) Panel 2012). Trends in outbreak data on FoNAO are, however, strongly influenced by very large outbreaks of considerable morbidity and mortality, such as the 2011 verocytotoxigenic *Escherichia*

MS 20141897 Submitted 14/11/2014, Accepted 13/2/2015. Authors Uyttendaele, Jacxsens, and Holvoet are with Dept. Food Safety & Food Quality, Ghent Univ., Ghent, Belgium. Author Jaykus is with Dept. of Food, Bioprocessing and Nutrition Sciences, North Carolina State Univ, Raleigh, N.C., U.S.A. Author Amoah is with Intl. Water Management Inst.—IWMI, Ghana. Authors Chiodini and Rao Jasti are with ILSI Europe, Intl. Life Sciences Inst., European Branch, 83 Ave. E. Mounier, B6, B-1200 Brussels, Belgium. Author Cunliffe is with Dept. of Health, Public Health, P.O. Box 6, Rundle Mall, 5000, South Australia. Author Korsten is with Dept. of Plant and Crop Sciences, Univ. of Pretoria, 0002, Pretoria, South Africa. Author Lau is with School of Chemical & Life Sciences, Nanyang Polytechnic, Singapore. Author McClure is with Mondelez Intl., Bayernwaldstrasse 8, 81737 München, Germany. Author Medema is with KWR, Watercycle Research Inst., Delft Univ. of Technology, Postbus 1072, 3430 BB, Nieuwegein, The Netherlands. Author Sampers is with Dept. of Industrial Biological Sciences, Ghent Univ. Campus Kortrijk, Kortrijk, Belgium. Corresponding author (E-mail: publications@ilsieurope.be).

coli (VTEC) O104 seed sprout outbreak in Germany. Even excluding this outbreak, FoNAO still caused 10% of outbreaks, 18% of cases, but only 8% of hospitalizations and 5% of deaths in the EU (EFSA Panel on Biological Hazards (BIOHAZ) 2012, 2013). In 2008, the World Health Organization (WHO) categorized leafy green vegetables as the highest priority in terms of fresh produce safety from a global perspective, as the most common produce items associated with outbreaks were greens-based salads and lettuce (World Health Organization 2008b).

Fresh produce can become contaminated with microbiological pathogens during production, at the processing/packing stage, and/or during preparation. Unfortunately, the importance of each of these different phases in the farm-to-fork continuum relative to pathogen contamination is unknown. However, it is clear that water is an important source of contamination, and over the years, there has been particular interest in the role of irrigation waters in this respect. Natural sources of water for irrigation include lakes and rivers, collected rainwater, desalinated sea water, and deep aquifers or shallow groundwater. The potential for microbial contamination of these water sources varies significantly depending on a variety of factors (Suslow and others 2003).

On an international scale, a particularly important factor when considering the quality of water used in fresh produce primary production is the availability of water resources, which is under increasing pressure. There are many reasons for this. For example, a growing population creates increased demand for water, particularly in urban areas. Climate variability causes greater unpredictability in precipitation, including periods of heavier rainfall as well as drought. Where there is dependence on groundwater, recharge often takes place at specific times of the year, so even relatively brief changes in rainfall patterns can have long-term effects. Even in parts of the world where water is often thought to be more plentiful (such as in India and South East Asia), there are significant pressures on water resources (Shah and others 2014).

In addition to increased water scarcity, human settlements produce a significant amount of wastewater that is rich in organic matter and which may be regarded as a resource if processed and disposed of properly. In fact, more and more regions are considering using this source of water as a valuable addition to the natural water resources available. Already wastewater reuse is practiced in many parts of the world and some countries have extensive experience with this technology (E.P.H.C. 2006). An important use for wastewater, after varying degrees of treatment, has been as a source of water for crop irrigation. This practice also reduces the impact of excess nutrients into surface waters and provides a source of plant nutrients in addition to the water needed for a wide variety of crops. This is practical particularly in water-stressed regions and where the source of wastewater is reasonably close to where the crops are grown.

Unfortunately, any water source can become contaminated with microbial pathogens. This is well known for drinking water and is the reason why efforts have been made over millennia to separate wastewater containing human and animal fecal matter from sources of drinking water. Consequently, a range of pathogens can also be present in waters used for fresh produce production and processing, and hence enter the food chain. Such hazards include both human-specific pathogens such as *Shigella* spp., norovirus, hepatitis A virus, *Cyclospora cayentanensis*, and zoonotic pathogens including verocytotoxin-producing *E. coli*, *Salmonella* spp., *Yersinia enterocolitica*, and *Cryptosporidium*. In addition, parasites such as tapeworms, that are of little consequence for drinking water (unless present inadvertently), are of major significance for food. While there are

some chemical hazards that are of significant concern for drinking water, such hazards are not generally an issue for foods grown using irrigation waters. This document focuses on microbiological issues. Its purpose is to discuss, in detail, the role of water quality in the safety of fresh produce. Accordingly, we describe the following areas:

- the epidemiological evidence supporting the role of water in primary production or at harvest in pathogen contamination and subsequent outbreaks of foodborne disease;
- the sources of water and methods used for irrigation during crop production and their impact on microbiological quality or potential for acting as a contributor to microbial contamination of fresh produce;
- the factors that impact survival of pathogens in water or in irrigated soil and fresh produce;
- control measures and guidelines for the management of water sources and water treatment, including microbiological criteria, to ensure the safety of fresh produce; and
- the role of testing and sampling to ensure appropriate water quality.

Fresh Produce and Microbial Food Safety Concerns and Water

Interest in the safety of fresh produce has grown almost exponentially over the last decade. Several comprehensive review articles have been produced, and the interested reader is referred to some selected ones for further details (De Roeber 1998; Sivapalasingam and others 2004; Johnston and others 2006; Hanning and others 2009; Leon and others 2009; Strawn and others 2011; Oliveira and others 2012). This section briefly describes produce-hazard pairs that have previously caused recognized outbreaks. The associated epidemiological investigations, with a focus on water serving as the source of contamination, will be described.

Pathogen contamination of fresh produce can occur at multiple locations across the farm-to-fork pathway. While water is an important vehicle for produce contamination, it is not the only one. With the exception of a few documented instances in which seeds have been contaminated with pathogens, the production phase is the earliest point at which fresh produce becomes contaminated with pathogens. This phase includes the steps of planting, growing, irrigating, and other activities and treatments associated with the production of the mature plant. Contamination of produce at the production phase frequently occurs as a consequence of exposure to contaminated water or soil. The former is of great interest to this report and will be discussed in detail throughout the article. Soil can be a source of contamination of crops if the production site was previously used for animal production and industrial dumping, or if biosolids/sludge, manure, or animal waste was applied as fertilizer or for waste disposal. Animal encroachment (birds, mammals, reptiles, and so on) is another important source of produce contamination preharvest, as is water runoff from surrounding areas where animal feces contaminates the land.

Outbreak investigations are an important and challenging component of epidemiology and public health and can prevent future problems by identifying contamination sources and mitigation strategies (Reingold 1998). Data availability on outbreaks associated with fresh produce, and in particular detailed information on outbreaks and sources of contamination, are diverse and scattered. Table 1 lists examples of commodities and pathogens that have been reported in various foodborne outbreaks linked to fresh produce consumption. The most common produce items associated

Table 1—Foodborne outbreaks linked to consumption of fresh produce.

Causative agent	Year	Country	Human cases	Implicated food	Country of origin	(Likely) source of contamination	Reference
VTEC O157	August–September 2013	UK	19 cases	Watercress	Domestic production	Wildlife entering the watercress farm or runoff water	(Public Health England 2014)
EHEC	June 2013	Sweden	19 cases	Fresh salad	Domestic	Irrigation water could be the likely source but not confirmed	(Edelstein and others 2013)
Salmonella Saintpaul	April–July 2008	USA	1500 cases	Jalapeno and Serrano peppers	Import from Mexico	The outbreak strain was isolated from 2 environmental samples, agricultural water, and Serrano peppers on a farm in Mexico which grew the peppers	(Behravesh and others 2011)
Salmonella Newport	2002 & 2005	USA	510 cases & 72 cases	Sliced tomatoes	Domestic production	The outbreak strain was traced back to farms on the Eastern Shore of Virginia, specifically to on-site ponds used for irrigation water	(Greene and others 2008)
Salmonella Thompson	October–December 2004	Norway (and probably larger EU outbreak)	21 cases	Rucola lettuce	Import from Italy	It is speculated that using water of nonpotable quality for irrigation of vegetables close up to harvest may lead to contamination of the products with a variety of pathogens	(Nygård and others 2008)
Salmonella Litchfield	2006/2007	Australia	26 cases	Papaya	Domestic	<i>Salmonella</i> Litchfield was not detected in papaya samples of the inspected farms; however, at one farm other serotypes of <i>Salmonella</i> were detected in untreated river water that was used for washing papaya	(Gibbs and others 2009)
<i>E. coli</i> O157	September–December 2006	USA	205 cases	Prepackaged spinach	domestic	The outbreak strain was isolated from one of the fields and in addition from river water, cattle faeces on a farm nearby and wild pig faeces. A potential cause was that the river functioned as a vector between the contaminated faeces and the irrigation wells used	(CFERT and others 2007)
<i>E. coli</i> O157	July–September 2005	Sweden	135 cases	Iceberg lettuce	domestic	The lettuce was irrigated by water from a small stream, and water samples were positive for Stx 2 by PCR. The identical VTEC O157 Stx 2 positive strain was isolated from the cases and in cattle at a farm upstream from the irrigation point	(Söderström and others 2008)
Norovirus and enterotoxigenic <i>E. coli</i>	January 2010	Denmark and Norway	260 cases	Lettuce (lollo bionda) used in sandwiches	Import from France	Unknown; mentioned that since neither norovirus nor ETEC are zoonotic agents, human fecal matter may have been the source of the contamination, possibly via contaminated water.	(Ethelberg and others 2010)
Norovirus	June–September 2005	Denmark	More than 1000 cases	Frozen raspberries	Import from Poland	Unknown; mentioned that contamination with norovirus may have occurred at farm level by fecally-contaminated irrigation water, during harvesting by infected farm workers and/or during processing and freezing by infected workers at company level.	(Falkenhorst and others 2005)
<i>Cyclospora cayentanensis</i>	May 2011	Canada	17 cases	Basil	Import from the USA	Unknown (usually contamination of the produce with sporulated oocysts through irrigation with contaminated water, or by spraying with pesticides or fungicides prepared using contaminated water. Other possibilities include handling of the produce by workers who were infected	(Hoang and others 2005)
<i>Cyclospora cayentanensis</i>	May–June 2009	Sweden	18 cases	Sugar snaps	Import from Guatemala		(Insulander and others 2010)
<i>Cyclospora cayentanensis</i>	May–June 1996	Canada and US	More than 1400 cases	Raspberries	Import from Guatemala		(Herwaldt and others Ackers 1997)
<i>Cyclospora cayentanensis</i>	2000	Germany	34 cases	Fresh green leafy herbs	Southern Europe	Probably fertilization with human waste or fecal contaminated water used to irrigate crops, prepare pesticides, or freshen or clean produce at their origin	(Doller and others 2000)
<i>Salmonella enterica</i>	November 1999–January 2000	Brazil	26 cases	Mango	Domestic	Mangoes imported from Peru which were due to exposure to untreated water (inadequately chlorinated water) in the final step of the fruit fly control program	(Beatty and others 2004; Sivapalasingam and others 2004)
<i>Cryptosporidium parvum</i>	October 1995	USA	31 cases	Apple cider	Domestic	The cider mill did not use drop apples, and apples were washed and brushed before pressing; however, cattle were present near the farm, and the apples were washed with water from a source later determined to contain <i>E. coli</i>	(Centers for Disease Control and Prevention 1997)

with these outbreaks have been leafy vegetables (spinach, lettuce, and lettuce mixes or salads), herbs, sprouted seeds, tomatoes, and berries. Cantaloupes, green onions, peppers, papaya, sugar snap peas, and a number of other commodities have also caused outbreaks. Just as there is a broad range of commodities associated with foodborne illness, the list of foodborne pathogens is also extensive. The biological hazards that dominate most reported produce-associated outbreaks are either zoonotic or human in origin. The most common zoonotic bacteria include *Salmonella enterica* (various serotypes) and verocytotoxin-producing *E. coli* O157, whereas outbreaks associated with *Campylobacter jejuni* and *Listeria monocytogenes* are relatively rare. Recently, however, *L. monocytogenes* has been associated with one of the deadliest produce-associated outbreaks, involving cantaloupe melons (Centers for Disease Control and Prevention [CDC] 2011). Human-specific bacterial pathogens such as *Shigella* spp. and other pathogenic *E. coli* (such as enterotoxigenic *E. coli*) are included. In addition, some of the human enteric viruses, particularly hepatitis A virus, human noroviruses, and rotavirus, have caused outbreaks, as have the parasitic protozoa *Cryptosporidium* and *Cyclospora*.

A portion of these foodborne illnesses associated with fresh produce originate from poor water quality used in the production or postharvesting washing. For example, irrigation pond water was responsible for a multistate tomato-associated *Salmonella* Newport outbreak in the United States (Greene and others 2008). Iceberg lettuce contaminated with *E. coli* O157 caused a large outbreak in Sweden, probably due to river water used for irrigation. The organism contained *vt2* genes, which on their own may not be responsible for the outbreak but they are together with other factors involved. However, the strain was only isolated from cattle upstream (Söderström and others 2008). Agricultural water was also the source of contamination in a nationwide 2008 US outbreak of *Salmonella* Saint Paul in peppers (Behravesh and others 2011). A more recent Enterohaemorrhagic *Escherichia coli* (EHEC) outbreak in June 2013 in Sweden was caused by fresh salad, components of which could have been contaminated by irrigation water, although this could not be confirmed (Edelstein and others 2013). Most recently (September 2013), a verocytotoxin-producing *E. coli* outbreak associated with the consumption of watercress was attributed to either wildlife entering the farm or contaminated runoff water (Public Health England 2014).

However, identification of the implicated food vehicle and/or the location of the point of food contamination in fresh produce-associated outbreaks are recurrent challenges. In 2012, in the EU, only 6.3% of 5363 outbreaks investigated had the same causative agent identified in the food vehicle or food chain and in human cases (EFSA 2014). Investigations of several *Cyclospora* outbreaks in June–August 2013, associated with the consumption of contaminated iceberg lettuce, were still unable to identify the causative agent (CDC 2013). But earlier reports from CDC in 2006 reported that potential water issues may have been related to the fresh spinach outbreak at that time which was attributed to surface runoff from grazing areas onto cultivated fields, construction of irrigation wells, depths to groundwater and groundwater–surface water interaction, and direct use of surface water for irrigation (CFERT and others 2007).

A major limitation of epidemiological investigation is that, in many instances, the true source of contamination is never ascertained and, in the absence of data, investigators can only speculate or assume a source. Such is the case for water; many outbreak investigations assume that the use of nonpotable irrigation water

just prior to harvest, or contaminated wash water, is responsible for pathogen contamination of produce. There is substantial danger in such assumptions, not just because they can be incorrect, but also because there is evidence that once a particular transmission pathway is identified, repeated outbreaks and investigations lead to a bias in causation (Lynch and others 2009).

Criteria for reuse of treated wastewater in irrigation are specific to a country or a region. In low-income countries, a range of alternative safety practices such as cessation of irrigation prior to harvesting, lowering of watering cans to reduce splashes from the soil, furrow irrigation, and so on are recommended to safeguard public health as much as possible in the local context (Keraita and others 2010; Amoah and others 2011). It is also important to note that the use of contaminated water in the dilution and subsequent application of fungicides and insecticides can also pose a significant microbial risk in a preharvest setting. Special attention to water quality should be given when using delivery techniques (for example, sprayers) that expose the edible portion of leafy vegetables directly to water, especially close to harvest time (Codex Alimentarius Commission 2003b).

Despite the vast amount of information in Table 1, these reported outbreaks are likely underestimates of the real situation. At national or international levels, an outbreak will receive widespread attention if the event (i) creates serious impacts on public health; (ii) is unusual or unexpected (the agent and/or produce type are unexpected, the circumstances of the outbreak are unique); and/or (iii) poses a significant risk of international spread with consequences for international travel or trade restrictions. The latter criterion is also the one that the WHO's INFOSAN alert system follows to identify potential international food-related events as threats to public health (World Health Organization 2008a; Rosenkötter and others 2014). In point of fact, many of the smaller outbreaks are never investigated. Finally, and perhaps most importantly, the vast majority of foodborne disease illness cases occur sporadically in the population, and these are not at all captured in routine epidemiological surveillance or outbreak investigations (Scallan and others 2011). Hence, Table 1 is in no way exhaustive, and much more information can be gleaned by consulting other sources of information. Such sources include the scientific literature, annual reports of national public health or food safety agencies (CDC or FDA in the United States; ECDC, EFSA, or RASFF in the EU, Food Standards Australia & New Zealand, and so on), and reports of outbreak investigations and epidemiological surveillance systems (such as Eurosurveillance, MMWR, and others). Local news media and dedicated Internet search engines (such as ProMedmail) are also information sources.

Most of the examples of fresh produce-associated outbreaks reported in Table 1 originate from Europe, North America, New Zealand, and Australia, as these locations have well-developed epidemiological surveillance systems; such systems are not available in much of the developing world. It is also important to note that, even for these countries, outbreak investigations may be biased or may differ geographically as a function of capacity, organizational structure, differences in trade flow, and so on. Outbreak data are rarely available from developing countries due to the lack of well-functioning surveillance and reporting systems. But it has been shown in some countries such as Senegal, South Africa, Mexico, and India that the quality of irrigation water and water for washing to maintain produce freshness influences microbial quality (presence of fecal indicator organisms and pathogens) (Minhas and others 2006; Ibenyassine and others 2007; Ndiaye and others 2011; Castro-Rosas and others 2012).

Table 2—Overview of agricultural practices in fresh produce production in different geographical areas.

Geographical area	Irrigation practice	Irrigation water	Cultivated fresh produce	Harvesting practices	Washing processes	Storage of produce	Reference
Sub-Saharan Africa	<ul style="list-style-type: none"> Watering cans and buckets Motorized pumps and hose Sprinkler Furrow Border or flood Drip irrigation Basin 	<ul style="list-style-type: none"> Rivers/streams Underground water Untreated wastewater (gray water) Underground water Rainwater stored in reservoirs 	<ul style="list-style-type: none"> Green leafy vegetables: e.g., lettuce, cabbage, spring onions Some fruit crops 	<ul style="list-style-type: none"> Mainly manual harvesting using knives, hand picking, cutlasses, etc. Mechanical harvesting (minimal) Special harvesting containers not used Harvesting implements mostly not cleaned 	<ul style="list-style-type: none"> Sometimes washed on farm with polluted irrigation water Clean water used for washing in markets 	<ul style="list-style-type: none"> Nonrefrigerated transport Storage at room temperature No storage facilities 	<ul style="list-style-type: none"> (Amoah and others 2007; Barno and others 2009; Ibenyassine and others 2007; Keraita and others 2007, 2010; Khalil and Comaa 2014; Mduli and others 2013; Yengoh and others 2010)
Middle East	<ul style="list-style-type: none"> Mostly surface irrigation Sprinkler Furrow Border or flood Drip (microirrigation) Basin 	<ul style="list-style-type: none"> Rivers/streams Underground water 	<ul style="list-style-type: none"> Green leafy vegetables Fruits Some tree crops 	<ul style="list-style-type: none"> Manual harvesting Mechanical harvesting 	<ul style="list-style-type: none"> Groundwater Potable water used for fresh-cut produce 	<ul style="list-style-type: none"> Refrigerated transport Storage at appropriate temperatures in cold storage room and refrigerator For non-RTE storage at ambient temperature 	<ul style="list-style-type: none"> (Bashour and others 2004; Feenstra and others 2000; Hussain and others 2002; Ongley 1996; Qadir 2008)
Central and South East Asia	<ul style="list-style-type: none"> Sprinkler Furrow Border or flood Drip (microirrigation) Basin Motor pumps 	<ul style="list-style-type: none"> Rivers/streams Underground water Untreated wastewater (gray water) Underground water 	<ul style="list-style-type: none"> Salads & green vegetables tomatoes fruits some tree crops 	<ul style="list-style-type: none"> Manual harvesting Mechanical harvesting 	<ul style="list-style-type: none"> Collected rainfall Tap water Groundwater Potable water used for fresh-cut produce Surface water (for first washing) Recycled surface water for transportation 	<ul style="list-style-type: none"> Refrigerated transport Storage at appropriate temperatures in cold storage room and refrigerator For non-RTE storage at ambient temperatures Ambient temperature for rural areas 	<ul style="list-style-type: none"> (Ahmad and Chua 2013; Basu and others 2012; Gorton and others 2011; Huong and others 2013)
Latin America	<ul style="list-style-type: none"> Sprinkler Furrow Border or flood Drip (microirrigation) Basin 	<ul style="list-style-type: none"> Rivers/streams Underground water Untreated wastewater Underground water 	<ul style="list-style-type: none"> Fruits and leafy vegetables Other tree crops 	<ul style="list-style-type: none"> Manual harvesting Mechanical harvesting (minimal) 	<ul style="list-style-type: none"> Collected rainfall Tap water Groundwater Potable water used for fresh-cut produce Surface water (for first washing) 	<ul style="list-style-type: none"> Refrigerated transport Storage at appropriate temperatures in cold storage room and refrigerator For non-RTE storage at ambient temperatures Ambient temperature for rural areas 	<ul style="list-style-type: none"> (Cardenas and others 2013; de Quadros Rodrigues and others 2014; Pereira and others 2002; Scott and others 2010)
OECD ^a countries	<ul style="list-style-type: none"> Sprinklers Drip systems Sheet irrigation Furrow Border strip Hydroponic mechanism 	<ul style="list-style-type: none"> Surface waters (rivers, reservoirs) Groundwater (wells, boreholes) Rainwater source for irrigation Tap water (to a lesser extent) 	<ul style="list-style-type: none"> Salads Green vegetables Fruit crops Many others Some tree crops 	<ul style="list-style-type: none"> Hand picking of fruits Mainly mechanical Machine harvesting 	<ul style="list-style-type: none"> Collected rainfall Tap water Groundwater Potable water used for fresh-cut produce Surface water (for first washing) 	<ul style="list-style-type: none"> Refrigerated transport Storage at appropriate temperatures in cold storage room and refrigerators 	<ul style="list-style-type: none"> (Steele and Odueru 2004; Tyrrel and others 2006)

^aOrganization for Economic Co-operation and Development.

Water Sources and Irrigation Methods Used in Fresh Produce Primary Production

Agricultural practices, including types of crops produced, water sources, and irrigation and harvesting practices, vary considerably around the world. Table 2 provides a summary of agricultural practices used in different geographical areas of the world. Clearly, there is considerable variability here. Fecally contaminated irrigation water is certainly a possible, and sometimes likely, source of pathogen contamination of fresh, ready-to-eat fruits and vegetables. Introduction of enteric pathogens from irrigation water is associated with either the source/type of water or the irrigation method (Brackett 1999; Steele and Odumeru 2004; Leifert and others 2008). Different sources and qualities of water are used for irrigation, with each of these having a different propensity to result in microbiological contamination of the crops. In addition, the method of irrigation plays an important role in the mode of contamination and transfer of bacteria, viruses, or protozoa to produce. In this section, we summarize key issues associated with water sources and irrigation methods applied at the fresh produce production phase of the farm-to-fork continuum.

Irrigation water sources

Water used for irrigation may originate from multiple sources: municipal water, rainwater, groundwater, surface water (open canals, impounded water such as ponds, reservoirs, and lakes), and wastewater (James 2006). The advantages and disadvantages of each are summarized in Table 3. Naturally, municipal water is of the best quality (but only available in some developed regions and quite expensive), followed by groundwater, rainwater, and surface water. The latter may or may not include discharges of treated (or untreated) human or industrial wastewater. Because of the acceptable quality and low cost of groundwater, this source of water is increasingly being used. However, the quality and sustainability of natural groundwater reservoirs is threatened in some regions by overabstraction. This results in the degradation of spring-fed rivers, destruction of wetlands, and chemical and microbiological contamination of the water (Krinner and others 1999; Reid and others 2003). In (semi) arid areas, desalinated seawater or brackish groundwater can also be used for irrigation purposes. Each of these water categories varies in microbiological quality as detailed below.

Rainwater or rain-harvested water is generally of relatively good microbial quality, albeit somewhat variable and of quality less than what is expected of potable water. The quality of rainwater depends in part on the means by which it is collected or transported. This can be illustrated with roof-harvested rainwater, which can be contaminated with pathogenic bacteria and protozoan parasites because of the presence of bird, insect, and animal droppings on roofs, especially immediately after relatively long periods of drought (Burch and Thomas 1998; Ahmed and others 2002). Water running off fields after heavy rainfall collects in lakes, rivers, or basins and can be heavily contaminated with pathogens from soil or fecal matter.

Groundwater (or borehole water) is generally of good microbial quality if infiltration of surface runoff is avoided (Burch and Thomas 1998). There can, however, be large variations between shallow groundwater and water from deeper aquifers. Although groundwater usually contains less organic matter than surface water, it may contain higher inorganic loads resulting in unpleasant colors and odors. The depletion rates of groundwater are accelerating worldwide, as evidenced by the fact that the rate at which humans are pumping dry the vast underground stores

of water has more than doubled since the early 1950s (Asano and Cotruva 2004; Foster and Kemper 2014). In general, borehole water shows less variability in terms of microbial load than rainwater (Steele and Odumeru 2004). Nonetheless, the potential for groundwater contamination from surface events, such as flooding or storm-related runoff from areas of concentrated manure accumulation, manure lagoons, or sewage treatment facilities, is well recognized (Oron and others 2001; Ibenyassine and others 2007; Rai and Tripathi 2007). There also are concerns based on well-water surveys and the prevalence of human illness derived from enteric virus contamination in this water (Gerba and Smith 2005; Pillai 1998). Thus, it is equally important to protect groundwater resources from microbial contamination sources.

Surface water includes lakes, rivers, creeks, ponds, and springs that come to the surface. Very often surface waters are contaminated due to discharges of (treated) wastewater, storm water runoff, livestock or wildlife feces, and so on. Also, surface waters show great variation in turbidity (Burch and Thomas 1998). More specifically, lakes tend to have better water quality than rivers, although lakes are also subject to surrounding sources of contamination from river inflow. Rivers, streams, and creeks have unpredictable water quality since activities upstream can rapidly change the levels of contaminants entering the flowing water. When surface water is used as the irrigation water source, drainage of contaminated water into the surface water reservoir can be avoided by constructing ditches, buffer strips, retention systems, and drainage systems. Potential overflow points should also be eliminated.

Seawater or brackish water, as with other surface waters, is subject to industrial and municipal waste discharges and river or stream runoff, possibly containing a wide range of human enteric pathogens. There are some crops having high salt tolerance, such as wheat and barley, and this property can be enhanced by selecting and breeding, thus providing crop varieties that can be irrigated with diluted seawater (Ghadiri and others 2006). However, in nearly all cases, seawater needs to be properly desalinated (such as seawater from which salt and other minerals are removed to a certain degree) by thermal processes or reverse osmosis before use in agriculture (Guler and others 2010). That process can achieve significant reduction of microorganisms. Although the costs of reverse osmosis membranes are high, the use of desalinated seawater might be economically feasible for high-value crops such as greenhouse vegetables and flowers (Yermiyahu and others 2007). Brackish groundwater (i.e., groundwater containing salt, but in lower concentration than seawater) can also be applied for irrigation when desalinated. However, the fact that groundwater is a limited resource, as opposed to seawater, should be taken into consideration (Muñoz and Fernández-Alba 2008).

It is generally believed that the use of untreated wastewater for irrigation presents significant health risks and, hence, is not a recommended practice (Pedrero and others 2010). Wastewater is usually of very poor physicochemical and microbiological quality and, consequently, requires intensive treatment prior to use in irrigation, unless other safety measures are in place when treatment is not feasible. Unfortunately, wastewater used for irrigation is often untreated or treated inadequately, particularly in developing countries (World Health Organization and others 2000). For example, it has been estimated that the percentage of effectively treated wastewater was 14% in Latin America and the Caribbean, 35% in Asia, 66% in Europe, and 90% in North America (Carr and Blumenthal 2004). Homsí (2000) estimated that only 10% of wastewater is treated in developing countries, resulting in about 20 million hectares of irrigation with insufficiently treated or

Table 3—Comparison between the different types of water sources used for irrigation.

Aspect	Municipal water	Groundwater	Collected rainfall water	Surface water
Definition (Codex Alimentarius Commission 2003c; Jacxsens 2010)	Water of potable quality offered by water companies	Water, seeped through from the surface and present in porous rocks below the surface, shallow wells or deep aquifers	Collected water from precipitation (rain, snow, ...)	Water from a source that is exposed to the environment like rivers/canals/lakes/open wells
Cost (example from one European country, i.e., Belgium)	Capacity compensation + approximately 3.3 euro/m ³ (www.pidpa.be)	First 499 m ³ are free of charge, between 500 and 30000 m ³ , one m ³ cost around 0.08 € (VMM 2013)	Free	Charging depending on the surface water if > 500 m ³ /y
Contamination sources	Pipelines, biofilm	Failing of septic systems, leaking sewer lines and from land discharge by passage through soils and fissures or interaction with surface water (Fong and others 2007; Hunt and others 2005; Lucena and others 2006; Steele and others Odumeru 2004).	Dust, organic matter, leaves, bird and animal excreta on the catchment areas (Evans and others 2006; Sazakli and others 2007).	Treated wastewater, discharge of raw sewage, municipal wastewater, storm-water runoff, runoff from urban and agricultural areas. Animals like birds, farm animals, and even humans are both indirect and direct contributors to the contamination (Geldreich 1991; Savichtcheva and others Okabe 2006; Sliva and others Dudley Williams 2001).
Weather impact	–	Heavy rainfall may lead to changes in the direction of water flow systems and flow through channels that would not normally occur which could lead to contamination (Hunter 2003).	Microbial profile found in rainwater systems was dependent on local environmental conditions and wind speeds/directions (Evans and others 2006). Rainfall after longer dry periods results in an increased presence of bacteria in the reservoirs (Schets and others 2005; Yaziz and others 1989). The first flush of rainwater carries most contaminants into storages (Yaziz and others 1989).	Storms, tides, or strong winds cause sediment resuspension, bacteria will also resuspend, resulting high bacteria levels in the water column (Ahn and others 2005; Bai and others Lung 2005; Parker and others 2010; Stumpf and others 2010). An additional increase in the numbers of organisms in the surface water is obtained due to heavy rainfall or storm flow through sewage overflow and surface runoff (Ahn and others 2005; Astrom and others 2009; Goyal and others 1977; Parker and others 2010; Rechenburg and others 2006).

diluted wastewater (Raschid-Sally and Jayakody 2009; Scott and others 2010). Municipal wastewaters generally contain pathogenic enteric bacteria, viruses, and intestinal parasites. Primary and secondary water treatment processes can eliminate 1 to 3 log₁₀ units of enteric microorganisms with an additional reduction 1 to 3 log₁₀ units achieved using tertiary treatments like filtration, all also depending upon the exact treatments used and the type of microorganism (bacteria, viruses, or protozoan (oo)cysts) (World Health Organization 2006). However, high microbial numbers might still be present in these purified wastewaters and additional disinfection practices should be applied if further elimination is required, as is frequently the case (Liberti and others 2000, 2001; Dell'Erba and others 2004; Koivunen and Heinonen-Tanski 2005; Falsanisi and others 2006). Treated wastewater is an increasingly relevant water source for irrigation, as it offers a year-round water supply and reduces the exploitation of natural sources, in particular the slow-recharging water layer (Lopez and others 2006). Minimum requirements of good practice to protect the health of the people using wastewater or excreta, or consuming products grown with wastewater or excreta, are provided by the WHO (World Health Organization 2006) and its more recent Guidance Notes (World Health Organization 2010).

Worldwide, however, most irrigation water derives from 2 main sources: surface water or groundwater reserves such as aquifers (Gleick 2000). In general, irrigation with surface water is expected to pose greater risk to human health than irrigation with water from deep aquifers drawn from properly constructed and protected wells, largely because of the ability to prevent

animal fecal contamination and runoff water from adjacent fields using the latter method (Suslow and others 2003). Most of these water sources are naturally replenished by precipitation. The exception to this is wastewater whose volume depends more on the population size contributing to the pool of wastewater. It is also important to note that different sources of water are very often mixed to obtain sufficient volume needed for certain water-intensive crop production settings and climatic conditions. Surely in times of water shortage, sources must be mixed, but the quality of the final water produced can vary and be unknown by the user. The identification of source water, combined with the definition of appropriate water quality, is vital to ensure the safety of irrigated products (Stine and others 2005).

Irrigation methods

Irrigation methods vary (usually by region) and each method may have its own potential to introduce human pathogens or, on occasion, even promote human pathogen growth on the product (Stine and others 2005). Irrigation methods range from very simple manual practices in the developing world to more sophisticated mechanical practices in the developed world. Commonly used irrigation methods include watering cans and buckets, motorized pumps with hosepipe (Obuobie and others 2010) (the latter are usually used in Africa and other developing countries), while sprinkler irrigation systems, irrigation by canals (furrows), drip irrigation, hydroponic cultivation, and so on, tend to be used in the developed world. Each irrigation method is discussed below in some detail.

Watering cans and buckets: small-scale farmers may use watering cans and buckets to fetch and manually carry water, from a water source, mostly shallow dug wells, streams, or dugouts, to the fields, followed by watering of crops through the spout or shower head of the can. This is, therefore, an overhead irrigation method. When men use this method, they usually carry 2 watering cans at a time, while the bucket system is mostly practiced by women and children (IPTRID 2001; Keraita and others 2002). Farmers using buckets and watering cans come in direct contact with water mainly by stepping in it while fetching it, or from water splashing on them while carrying it and during watering; highly contaminated water can present a health risk to the farmers themselves. If the water is contaminated with microbial pathogens, the likelihood for subsequent crop contamination is very high because of the combination of an overhead application and a large surface area.

Other *surface irrigation methods* include *flood irrigation* (water applied over the entire field to infiltrate the soil); *canal or furrow irrigation* (water applied between ridges, for example, level and graded furrows, contour furrows, corrugations, and so on); and *sprinkler irrigation* (in which water is applied in the form of a spray and reaches the soil more or less like rain, from travel sprinklers, spray guns, portable and solid-set sprinklers, and so on). The flood irrigation system results in complete coverage of the soil surface with water and is normally not an efficient irrigation method. This system can also result in contamination of root crops or vegetable crops growing near the ground. Because it results in direct farm worker exposure, more so than any other method, flood irrigation poses the greatest health risks to both farmers and consumers when contaminated irrigation water is used. Similarly, sprinkler irrigation facilitates the contamination of ground crops (exposing the edible portion of the produce directly to water, a particular problem if applied close to harvest time), fruit trees, and farm workers. Splashing of sprayers can create recontamination of the crop surface from the soil (Marites and others 2010). In addition, pathogens contained in aerosolized effluent may be transported downwind and create a health risk to nearby residents (Fattal and others 1987). Risk associated with spray irrigation may increase if the irrigation event occurs immediately after a high wind lapse (Barker-Reid and others 2009).

In *subsurface irrigation*, water is supplied through deep surface canals or buried pipes beneath the root zone in such a way that it wets the root zone by capillary action, whereas in *drip irrigation* water is applied around each plant or a group of plants so as to wet only the root zone and to limit the moisture to a relatively local application. Relatively speaking, these methods certainly provide the greatest degree of health protection for farm workers and consumers, especially if the methods are automated. Drip irrigation and well-maintained furrow irrigation also limit contamination of leaf surfaces (Qadir 2008). Plants grown without soil, such as in hydroponic systems, absorb nutrients and water at varying rates, constantly changing the composition of the recirculated nutrient solution. Hence, water used in hydroponic culture should be changed frequently or, if recycled, a water treatment method should be applied.

The method of irrigation plays an important role in the transfer of contamination to crops. It is important to note that irrigation distribution networks are designed to meet peak demands, which might create, in some parts of the network, low-flow conditions that can contribute to the deterioration of microbial quality of water. Also, maintenance of the water delivery systems is important as biofilms can increase the contamination between the source and the tap (Szewzyk and others 2000; Hallam and others 2001).

The most often applied systems of irrigation in professional crop production and the advantages and disadvantages associated with these different irrigation methods are summarized in Table 4. It is clear that contamination and transfer of pathogens depends on the irrigation method and on the nature of the produce (SCF 2002). Subsurface or drip irrigation lowers the risk of transfer to growing plants compared to furrow and sprinkler irrigation, by minimizing the exposure of the irrigated water to the produce (Oron and others 2001; Enriquez and others 2003; Hamilton and others 2006; Song and others 2006). Furthermore, subsurface or drip irrigation lowers the risk of splashing of contaminated soil on vegetables (Ntahimpera and others 1999; Pietravalle and others 2001; Girardin and others 2005; Franz and others 2008; Cevallos-Cevallos and others 2012).

Pathogen contamination by irrigation water is of greatest concern when irrigation is done right before harvest. For example, water containing 2.5 log CFU *Salmonella* spp. was sufficient for contamination and persistence of the pathogen on plants for at least 48 h after spray irrigation (Kisluk and Yaron 2012). Other studies have reported *E. coli* persistence after spray irrigation for up to 27 days (Erickson and others 2010). Hence, unless the water quality is well controlled and potable, spray irrigation is best applied in the early stages of plant growth, thus maximizing the opportunity for pathogen die-off.

Based on the above reflections, Table 5 demonstrates the risk ranking of lower to higher risk of combinations of sources of water with different irrigation methods and types of crop. The highest risk of contamination can be attributed to the combination of raw/poorly treated wastewater to be used for surface irrigation with watering cans as applied to low-foliar plants such as lettuce or root crops such as onions or carrots. Ultimately, however, the type of irrigation method chosen by a grower depends on several issues, including the groundwater depth, types of water sources available, local cost of these water sources, cost of irrigation equipment/infrastructure, soil type and slope, and crop type or applicability of crop rotation(s) (Mena 2006).

The Behavior of Microbial Hazards in the Production Environment

In considering microbial pathogen contamination of fresh produce, it is important to understand that, once the microbes are introduced into water and via water into soil or plants, the factors impacting their ability to survive, and perhaps even grow, under given climatic and environmental conditions or stages of crop production are important. The ability of pathogenic organisms to attach, survive, and grow on the surface of various fruit and vegetables is dependent upon (i) the metabolic capabilities of the pathogens themselves; (ii) the unique set of intrinsic factors possessed by a particular produce item; and (iii) the extrinsic ecological factors that naturally occur in or on the produce at various stages of production, processing, distribution, and/or preparation (Beuchat 2002). The survival of pathogens is important as it can impact the likelihood of an outbreak (Fonseca and others 2011). In general, the survival of pathogens in pristine water decreases with increasing temperatures (Rhodes and Kator 1988; González and Hänninen 2012). However, increasing nutrients and high organic load can also increase survival. For example, the viable counts of *Campylobacter* spp. decreased below detection limits within 5 days at 25 °C and within a maximum of 70 days at 4 °C (González and Hänninen 2012). Thomas and others (2002) found 18 times higher decay rates for *Campylobacter* spp. at 20 °C. The survival of *E. coli* O157 in surface water strongly decreased with increasing temperatures; it survived 8 weeks at 25 °C compared to 13 weeks at 8 °C

Table 4—Comparison of the (dis)advantages of the different irrigation methods.

Aspect	Canal/furrow irrigation	Sprinkler/overhead irrigation	Drip/subsurface irrigation
Definition (Eurostat 2003)	Leading of water along the ground, either by flooding the whole area or leading the water along small furrows between the crop rows, using gravity as a force	Irrigating the plants by propelling water under high pressure as rain over the parcels	Irrigating the plants by placing water low by the plants drop by drop or with microsprinklers or by forming fog-like conditions
Advantages (Ghassemi and others 1995; Verbeten 1998)	Low capital costs	Suited for a wide range of slopes, soils, and crops	Increased uniformity, soil structure is preserved, water is saved because of reduced evaporation and a correct control of water quantities and nutrients reaching plants is possible
Disadvantages (Ghassemi and others 1995; Verbeten 1998)	Uneven penetration of the water	Avoidance of uneven penetration of water and its subsequent waste	Correct control of water quantities and nutrients reaching plants is possible
	Water application onto the field may be uncontrolled Not suited for all slopes and soils	High initial cost of equipment The higher operation costs compared with surface irrigation The need of a pumping plant and the requirement of energy	High capital costs Obstruction of small drippers because of water impurities Creation of an area of permanently saturated or near-saturated soil favoring the development of plant or animal pests

Table 5—Levels of risk associated with different source waters, irrigation methods and crop types.

Level of risk	Source water ^a	Irrigation method	Crop type
Lower	<ul style="list-style-type: none"> • Municipal potable water • Groundwater collected from deep wells/bores • Rainwater (collected in closed systems) • Groundwater from shallow wells/bores 	<ul style="list-style-type: none"> • Subsurface • Drip • Furrow 	<ul style="list-style-type: none"> • Root crops (e.g., onions) • Low foliar (e.g., lettuce) • Off ground (e.g., tomatoes)
Higher	Adequately treated wastewater Rainwater (collected in closed systems) Surface waters in proximity to animals/human habitation Raw/poorly treated wastewater	<ul style="list-style-type: none"> • Spray • Surface irrigation with watering cans 	<ul style="list-style-type: none"> • Fruit trees (e.g., apple, mango) • Low foliar (e.g., lettuce)/root crops (e.g., onions)

^aFrom FAO/WHO Microbiological hazards in fresh leafy vegetables and herbs 2008.

(Wang and Doyle 1998). *Salmonella* spp. survived 24 weeks in fresh-water microcosms at ambient temperature (30 °C) compared to 58 weeks at temperatures of 5 °C (Sugumar and Mariappan 2003).

After irrigation, the ability of enteric bacteria to survive in the hostile environment of the phyllosphere is debatable. Stress conditions on plant surfaces can restrict pathogenic bacterial survival (Brandl 2006; Warriner and Namvar 2010). Enteropathogens can adapt to the phyllosphere environment but may fail to compete with indigenous epiphytes (Janisiewicz and others 1999; Brandl and Mandrell 2002; Cooley and others 2006; Warriner and Namvar 2010). Between 30% and 80% of the total bacterial population on a leaf surface is located in biofilms having an increased survival rate (Morris and Monier 2003). Even if human pathogens cannot produce homogeneous biofilms, they may become entrapped in heterogeneous biofilms produced by nonpathogenic bacteria, making them much more restive to stress conditions (Fett 2000). However, it has been reported that *E. coli* O157:H7 may not preferentially colonize biofilms produced by natural microbiota on lettuce leaves (Seo and Frank 1999).

A number of key factors are likely to influence bacterial death on the phylloplane, the most important being low humidity, high temperatures, exposure to UV, and wind-mediated drying of the leaf surface (Gras and others 1994; Hutchison and others 2008; Moyne and others 2011; Oliveira and others 2012). The survival and growth of certain enteric pathogens on plants depends on the relative humidity (RH). Low RH has been proposed as one of the main factors limiting survival of bacteria on plant surfaces

(Medina and others 2012; Oliveira and others 2012). For instance, *Salmonella* spp. populations declined rapidly under low RH on cilantro, whereas the organisms were able to grow on cilantro leaves under humid conditions. Phylloplane bacteria are also efficient in UV-induced DNA damage repair (Heaton and Jones 2008). Enteropathogens encounter osmotic stress when passing through the host gut, which may induce cross-resistance to stresses encountered on the leaf (Brandl 2006). Protection from environmental stresses may be facilitated by movement into the internal tissue of the plant. Enteropathogens in irrigation water can be taken up by the root system, or via wounds or other structures such as stomata, and enter the edible portion of the plant (Janisiewicz and others 1999; Seo and Frank 1999; Solomon and others 2002; Zhang and others 2009). However, despite a lower survival of pathogens in the field in the warmer seasons, there are higher chances of pathogen introduction at these times (Fonseca and others 2011).

Several studies have examined the persistence and survival of pathogens on lettuce through the application of irrigation water, manure, or direct inoculation of lettuce with soil and manure. Nevertheless, the comparison of data from individual studies is difficult due to variability in experimental design and conditions, plant species, cultivars, maturity at inoculation, bacterial strains and their cultivation, and analytical methods (Delaquis and others 2007). A summary of individual studies carried out with leafy vegetables is provided in Tables 6 and 7. The tables emphasize the big differences in experimental design between studies. Sometimes artificially high inoculation levels (5–9 log₁₀ CFU/g or mL)

Table 6—Irrigation with contaminated water on leafy vegetables and the subsequent survival of enteric bacteria in lettuce and soil.

Setting	Produce	Bacteria	Inoculum (log CFU/mL)	Irrigation method	Survival in soil after inoculation	Survival on produce after inoculation	Reference	
Field	Leafy green lettuce	<i>E. coli</i> O157	8	Spray	ND	27 days	(Erickson and others 2010)	
	Iceberg lettuce	<i>E. coli</i>	8–9	Spray	7 days	1 day	(Fonseca and others 2011)	
	Iceberg lettuce	<i>E. coli</i>	8–9	Drip	7 days	< 1 day		
	Iceberg lettuce	<i>E. coli</i>	8–9	Furrow	15 days	< 1 day		
	<i>Lactuca sativa</i> L.	<i>E. coli</i> O157:H7	5	Spray	140 days	56 days	(Islam and others 2004)	
	Parsley	<i>Salmonella enterica</i>	8.5	Spray	ND	4 weeks	(Kisluk and others Yaon 2012)	
	<i>Lactuca sativa</i> L.	<i>E. coli</i> O157:H7	4	Surface	ND	15 days	(Mootian and others 2009)	
	Lettuce, parsley, tomato, and pimento	Untreated wastewater	ND	Surface	ND	3 days	(Melloul and others 2001)	
	Lab	<i>Lactuca sativa</i> var. <i>longifolia</i>	<i>L. innocua</i>	7	Spray	ND	4 weeks	(Oliveira and others 2011)
		<i>Lactuca sativa</i> var. <i>Longifolia</i>	<i>E. coli</i> O157:H7	7	Spray	ND	4 weeks	(Oliveira and others 2012)
Green ice lettuce		<i>E. coli</i> O157:H7	7	Surface	ND	20 days (6/32 samples positive)	(Solomon and others 2002)	
Green ice lettuce		<i>E. coli</i> O157:H7	7	Spray irrigation	ND	20 days (29/32 samples positive)		
Butterhead lettuce		<i>E. coli</i> O157:H7	4	Spray	ND	30 days	(Solomon and others 2003)	
Butterhead lettuce		<i>E. coli</i> O157:H7	2	Spray	ND	15 days		
Spinach	<i>E. coli</i> O157:H7	5	Spray	ND	6 days	(Wood and others 2010)		

were used, such as in experiments to investigate the use of contaminated compost and irrigation water on the ability for cross-contamination, survival, and internalization of human pathogens in soil or lettuce. Still, the survival of pathogens after application of irrigation water or manure ranged from 1 day up to 2 months on lettuce and more than 7 months in soil depending on inoculation level and season (Hutchison and others 2005; Liu and others 2013). Survival of foodborne pathogens on produce is significantly enhanced once the protective epidermal barrier has been broken either by physical damage, such as punctures or bruising, or by degradation by plant pathogens or spoilage organisms (bacteria or fungi). These conditions can also promote the multiplication of human pathogens, especially at ambient temperatures. Injured cells and released cell fluids provide a nourishing environment for microbial growth. Certain crop management practices and/or extreme weather conditions (such as heavy rain, hail, or strong winds) might influence tissue susceptibility for contamination and internalization with foodborne pathogens by affecting plant physiology, tissue structure, and microbial ecology.

Prevention and Control Measures for Irrigation Water Quality

To ensure the safety of fresh produce, and simultaneously safeguard the health of crop producers and their staff, a set of guidelines, namely, Good Agricultural Practices (GAPs), have been released. These good practices are defined at the international level in the “Codex General Principles on Food Hygiene” (Codex Alimentarius Commission 2003b), with guidance specific to fresh produce production further developed in “CAC/RCP 53–2003 Code of practice for fresh fruits and vegetables.” This particular Codex document provides explanations of good practices to minimize the contamination of fresh produce during cultivation and (post)-harvest practices. There are many suggested practices, but, because of the importance of water as a potential source of contamination, significant parts of GAP documents focus on water. The Codex Committee on Food Hygiene guidelines for control of virus contamination of food (Codex Alimentarius Commission 2012) also recommends that efforts should be made to use only potable or clean water (this is water quality that does not affect the wholesomeness of the food) during production. In parallel to the Codex Alimentarius documents, several guidelines and quality assurance standards were developed for the primary production of fresh produce on the initiative of national competent authorities, fresh produce industry associations, or as voluntary private standards and marketing agreements in the fresh produce supply chain. For example, in the United States, there are general and specific guidance documents provided by the U.S. FDA (FDA 1998, 2009a, 2009b, 2009c, 2009d). Guidance is also provided by specific commodity groups (United Fresh Produce Association 2002, 2005, 2008 2010). GlobalGap is the European retailers private collective standard set and also acts as an organization for benchmarking other voluntary standards in agricultural production (including fresh produce production) around the globe enabling certification of GAPs (www.globalgap.org). An alternative organization, SQF (Safe Quality Foods), initially developed in Australia in the early 1990s and currently owned and managed by the Food Marketing Institute in the United States, has elaborated the SQF 1000 Code for primary production as a quality assurance standard for certification of GAPs (www.sqfi.com). Another quality assurance standard developed on a national level is *Integrated Chain Quality Management* (ICQM) in Belgium (www.vegaplan.be). The ICQM Standard applies specifically

Table 7–Inoculation of soil or leafy greens and the subsequent survival of enteric bacteria.

Setting	Produce	Bacteria	Inoculation Inoculation	Survival in soil after inoculation	Survival on produce after inoculation	Reference
Field	<i>Lactuca sativa</i>	<i>E. coli</i> O157:H7	Composts: 7	154 days	77 days	(Islam and others 2004)
	<i>Lactuca sativa</i> L.	<i>E. coli</i> O157:H7	Soil and manure-amended soil: 1, 2, 3, and 4	ND	15 days	(Mootian and others 2009)
Lab	Romain lettuce	<i>E. coli</i> O157:H7	Plants: 5, 4–6, 4	Soil: up to 15 days	Lettuce: up to 35 days	(Moyné and others 2011)
	Spinach	<i>Salmonella enterica</i>	Soil: 4, 7	ND		(Arthurson and others 2011)
	Romain lettuce	<i>E. coli</i> O157:H7	Soil: 6	ND	All samples after 7 days 40% after 14 days 20% after 21 days	(Theofel and others 2009)
	Romain lettuce	<i>E. coli</i> O157:H7	Lettuce leaves 6.5 log CFU/leaf 24 h, 23°C, <50% humidity	ND	Reductions of 1.8 to 3.3 log CFU/leaf	(Theofel and others 2009)
	Romain lettuce	<i>E. coli</i> O157:H7	Fresh-cut lettuce 3.5 5 days, 5°C and 20°C, <50% humidity	ND	Increase at 20°C after 24 h decrease at 5°C after 1, 2, and 5 days	(Theofel and others 2009)
	Romain lettuce	<i>E. coli</i> O157:H7 and <i>S. enterica</i>	Plants: 4 3 days, 28°C, 100% relative humidity	ND	100-fold increase <i>E. coli</i> O157:H7 155-fold increase <i>S. enterica</i>	(Brandl and others Amundson 2008)
	Romain lettuce	<i>E. coli</i> O157:H7 and <i>S. enterica</i>	Harvested leaves: 4 3 days, 28°C, 100% relative humidity	ND	500-fold increase <i>E. coli</i> O157:H7 and 740-fold increase <i>S. enterica</i>	(Brandl and others Amundson 2008)
	Romain lettuce	<i>E. coli</i> O157:H7	Soil: 5 20°C, 70% relative humidity	ND	36 days	(Ibekwe and others 2006)
	<i>Lactuca sativa</i> L cv. Dublin	<i>E. coli</i> O157:H7, <i>Salmonella</i> serovar Typhimurium	Manure: approximately 7	<i>E. coli</i> O157:H7 up to 56 days <i>Salmonella</i> Typhimurium > 56 days	One lettuce root was positive lettuce <i>E. coli</i> O157	(Franz and others 2005)
Green-house	NA	<i>E. coli</i> O157:H7	Inoculated manure mixed with unautoclaved or autoclaved soil: 6–7	Autoclaved soil: 231 days 21°C	ND	(Jiang and others 2011)
	Crisphead Lettuce	<i>E. coli</i> O157:H7	Bovine manure: 4	8 weeks	Not detected	(Johannessen and others 2005)

Table 8–Water sources and quality guidelines proposed in different manuals on good practices in fresh produce primary production.

Criteria	Codex Fresh produce, (Codex Alimentarius Commission 2003a)	GlobalGap (Global Gap 2013)	SQF 1000 (SQF Institute 2010)	McDonald's Worldwide Quality Systems 2011)	FSMA (New England Farmers Union 2013)	ICQM (Primary Production 2013)	Spain (BOE 2007)	Norway (Matmerk KSL 2010)
Sources of irrigation water	Identify the sources of water, assess its microbial quality, and its suitability for intended use	No untreated sewage water	A known clean source or treated to make it suitable for use	Adequate distances between water sources and potential sources of contamination must be maintained. Water sourced from a well or bore hole must not be closer than 200 feet (60 m) to areas of untreated manure accumulation. Water sourced from open surface water must not be closer than 100 feet (30 m) to areas of untreated manure accumulation for sandy soils and no closer than 200 feet (60 m) for loamy or clay soils. Reclaimed recycled water is not permitted unless it meets (treated or untreated) certain criteria	Adequate for its intended use	Creek, open well, drilled well, rain, potable or vegetable wash water, or water used in the processing of vegetables such as blanching, sterilizing	Microbial criteria have only been established for the use of treated wastewater for the irrigation of crops that are likely to be eaten uncooked	The irrigation water source should be protected against contamination
Microbio-logical standards	–	If treated sewage water is used, water quality complies with WHO guidelines.	Based on the hazard analysis, best practices within country of production and any application legislation, if applicable	The geometric mean of generic <i>E. coli</i> of the 5 most recent samples must be less than 126 MPN/100 mL, with no single sample > 235 MPN per 100 mL	Water that may come in contact with the harvestable portion of produce must meet a standard of no more than 235 CFUs of generic <i>E. coli</i> per 100 mL throughout the growing season	–	<i>E. coli</i> : ≤100 CFU/100 mL, sampling plan 3 classes: n = 10; m = 100 CFU/100 mL; M = 1000 CFU/100 mL; c = 3 Intestinal nematode: ≤1/10 L	Water quality has to be "close to drinking water quality" (not specified) and the last day of irrigation before harvesting should be documented
Frequency of testing	Depend on the water source and the risks of environmental contamination	A frequency according to the results of the risk assessment every year	Decided by the hazard analysis, best practices within country of production and any application legislation	A set of 5 samples must be collected prior to harvest. Samples taken must be at least 18 h apart and not longer than 30 days since the last sample was taken	River or natural lake water: every 7 days during growing season, water reservoir from groundwater: once a month, groundwater: at the beginning of the season and every 3 months thereafter	–	–	At least one water sample should be analyzed each year

to agricultural crop production and horticulture and describes the minimum requirements for producers and workers on good practices to gain access to the high-value fresh produce market. In Norway, KSL Matmerk is a private initiative providing guidance and a quality system for agriculture. McDonald's Corporation has issued its own rigorous food safety standard for fresh produce production, and there are many other private collective or individual company-based standards and quality assurance programs available. Some of the guidelines on use of water sources and prerequisites on water quality and sampling and testing are shown in Table 8.

With regard to the legal framework demanding implementation of GAPs, in Europe the EC Regulation No 852/2004 (European Commission, 2004) on the hygiene of foodstuffs lays down general hygiene requirements to be respected by food businesses at all stages of the food chain, including at primary production. Thus, there is a legal obligation to comply with requirements for good hygiene practice and thus to prevent the contamination of food of plant origin also at primary production. Some European countries also have explicit legislation referring to the quality of water to be used in primary production (for example, Spain) (Table 8). In association with the U.S. FDA Food Safety Modernization Act (FSMA), regulations that result in mandatory GAPs adherence will inevitably be instituted, although these remain in developmental stages at the time of this writing.

Overall, several preventive control measures can be practiced on the farm in an effort to avoid microbial contamination of irrigation water. Although it is impossible to completely prevent and, if occurring, eliminate such contamination, careful attention to controls can minimize risk. For a further review of the most effective preventive measures and interventions, one is referred to the above-mentioned Codex Alimentarius Commission documents, the opinions issued in 2014 by EFSA on the risk posed by pathogens in food of nonanimal origin (EFSA 2014) or the review by Gil and others (2015).

As mentioned in many Codes of Practice for primary production of fresh produce, the importance of selecting a high-quality irrigation water source cannot be overemphasized. It is essential that, on a regular basis, sanitary surveys of water reservoirs and distribution systems are executed. These should focus on the integrity of surrounding protective structures, identifying potential point source and nonpoint source confluences (such as drainage into these systems). If the evaluation concludes that (human or animal) fecal contamination of the water in a specific area is at levels that may compromise the quality of the water and thus the safety of crops, appropriate interventions should be taken. The most important intervention used to address pathogen risks in irrigation water, if judged to be of insufficient quality, is water treatment. Treatment methods correspond approximately to what is used for sewage water treatment and include coagulation, flocculation, filtration, and disinfection. Solar irradiation is also suggested to reduce the levels of pathogenic microorganisms in irrigation water. Other options that have been considered to improve the microbial quality of surface waters include sand filtration or storage in catchments or reservoirs to achieve partial biological treatment before use (Carr and Blumenthal 2004). Overall, it is recommended to use a disinfection treatment if using water from open reservoirs that are prone to human or animal fecal contamination (and thus likely pathogen contamination). Disinfectant treatments of surface or well water include chlorination, use of peroxyacetic acid, and UV treatment. Ozonation has also been described as a possible disinfection treatment for irrigation water (Suslow and others 2003). It can be difficult to choose the technology that is the best fit for a

specific situation, as the performance of the water treatment process will depend upon physicochemical and microbial parameters associated with the water to be treated. Selection of technology will also relate to aspects including capital and operational costs, complexity of the technology, required monitoring, and safety issues (Van Haute and others 2013).

The Role of Testing and Monitoring in Ensuring Safe Water in Fresh Produce Production

Some Codes of Practice demand growers to have the water they use periodically tested for microbial contaminants. Depending on the type of water source and method of irrigation, microbial sampling may be recommended at different frequencies to verify the functionality of good agricultural practices. Testing is costly; if testing is applied, it is important that an agreement is made on the frequency and location of sampling; the sampling method and volume; the microbial parameters to be analyzed; the method of detection or enumeration of these microbial parameters; the interpretation of test results, including specifications and/or microbiological criteria; and the types of actions to be taken upon noncompliance. At present, there is no widespread agreement regarding the microbiological guidelines for irrigation water. In most cases, actual pathogen contamination of waters and fresh produce is probably quite rare, particularly in the developed world. Furthermore, direct pathogen screening is expensive, time-consuming, and difficult to interpret (Savichtcheva and Okabe 2006). Pathogens tend to be nonuniformly distributed in water, which complicates the interpretation of negative test results. In most cases, generic *E. coli* are used as indicator organisms, as their presence relates to fecal (animal or human) pollution. Alternatively, fecal coliforms may be used for this purpose. Total coliforms can be analyzed to indicate failures in control measures. As operational indicators, total coliforms may provide information on the adequacy of water treatment and on the microbial condition of the water distribution system at the point of application. It should be noted, however, that the presence of total coliforms in the (tank) water or the distribution system, without further discrimination, is of no immediate public health significance. Nonetheless, the presence of coliforms is still an indicator of inattention to “best practices” and should prompt further actions, such as sanitary survey of the construction of the water network, the input to the tank water, control of the water treatment, storage conditions, potential for regrowth of microorganisms, and so on.

As the pathogens associated with fresh produce outbreaks are almost always of fecal origin, a good indicator should correlate with the presence of fecal contamination. Historically, a subset of the coliforms, the fecal coliforms (those coliforms which ferment lactose with the production of acid and gas within 48 h at 44.5–45.5 °C in EC broth) have been most widely used in sampling and testing of water quality for this purpose. The major genera represented in the fecal coliform group are *Enterobacter*, *Citrobacter*, and *Klebsiella*, which are not always of fecal origin, although the majority of the fecal coliforms are strains of *E. coli*. A WHO world survey indicated that most European rivers contain mean fecal coliform counts of 1000 to 10000 per 100 mL (World Health Organization 1989). There are, in some countries or states, guidelines on appropriate microbial quality specifications for surface water or (treated) wastewater to be used for irrigation based on testing for fecal coliforms (Table 9). Guidelines for microbial water quality of surface water for irrigation are usually less stringent than those of wastewater for unrestricted irrigation due to the assumption that enteric viruses and other human pathogens are present in lower

Table 9—Irrigation water quality guidelines and regulations.

Country/region	Water type	Regulation/guideline	Criteria ^{a,b}	Reference
Australia New Zealand ^c	Irrigation water for nonfood crops (trees/flowers): Secondary treatment or primary treatment with lagoon detention	Guideline	< 1000 <i>E. coli</i> per 100 mL	
Australia New Zealand	Irrigation water for commercial crops raw or unprocessed (salads crops and spray irrigation): Advanced treatment to achieve total pathogen removal required (e.g., secondary, filtration, and disinfection)	Guideline	< 1 <i>E. coli</i> per 100 mL	
Australia New Zealand ^{e,f}	Irrigation water for commercial food crops: Secondary treatment with >25 days' lagoon detention and disinfection	Guideline	< 100 <i>E. coli</i> per 100 mL	
Canada	All	Guideline	1000 total coliforms per 100 mL < 100 fecal coliforms per 100 mL	(Steele and others Odumeru 2004)
Canada (Alberta)	Surface water	Guideline	1000 total coliforms per 100 mL < 100 fecal coliforms per 100 mL	(Steele and others Odumeru 2004)
Canada (British Columbia)	All	Guideline	200 fecal coliforms per 100 mL 77 <i>E. coli</i> per 100 mL: < 20 fecal streptococci per 100 mL	(Steele and others Odumeru 2004)
Canada (Saskatchewan)	Surface water	Guideline	1000 total coliforms per 100 mL < 100 fecal coliforms per 100 mL	(Steele and others Odumeru 2004)
Italy	(Treated) Wastewater	Regulation	10 <i>E. coli</i> per 100 mL <i>Salmonellae</i> absent in 100 mL	(Cirelli and others 2008)
Spain ^c	(Treated) Wastewater	Regulation	100 <i>E. coli</i> per 100 mL < 1 nematode egg in 10 L	(BOE 2007)
USA	Surface water	Guideline	< 126 <i>E. coli</i> per 100 mL	(LGMA 2012; US Environmental Protection Agency 2003)
USA	(Treated) Wastewater	Guideline	Fecal coliforms absent per 100 mL	(US Environmental Protection Agency 2004)
California (USA)	?	Regulation	< 2.2 total coliforms per 100 mL Fecal coliforms absent	(Steele and others Odumeru 2004)
WHO	Wastewater	Guideline	< 1000 fecal coliforms per 100 mL < 1 nematode egg per L	(Blumenthal and others 2000; World Health Organization 2006)

^aAll values per 100 mL, unless otherwise stated, TC = total coliforms, FC = fecal coliforms, EC = *E. coli*. ^bSpecifics of sample value calculation, such as geometric mean, minimal number of samples, period of sampling, percentage of samples that may deviated from the target value, etc. are not mentioned here. ^cDirect contact of irrigation water with edible parts. ^dNo direct contact of irrigation water with edible parts. ^eCrops with limited or no ground contact and eaten raw (e.g., tomatoes, capsicums)—drip irrigation and no harvest of wet or dropped produce. ^fCrops with ground contact with skins removed before consumption (e.g., watermelons)—if spray irrigation, minimum 2 days between final irrigation and harvest.

concentrations in surface water (Gerba and Choi 2006); or in the context of microbiological criteria, fecal coliforms in surface water may originate from sources other than sewage or waste effluents, which is certainly the case in hot climates.

Actually, with better detection methods available, *E. coli* is now the indicator of choice for fecal contamination originating from warm-blooded animals (including humans) (Mossel 1978, 1983). The presence of generic *E. coli* provides evidence of an increased

likelihood of potential contamination of food or water by ecologically closely related pathogens. Holvoet and others (2014) showed that the use of water with *E. coli* levels higher than 2 log₁₀/100 mL needs to be avoided, as 42% of the water samples with values exceeding this contained a pathogen (*Salmonella* or *Campylobacter* isolates) or the presence of verocytotoxin genes (indicative of the presence of pathogenic *E. coli* genotypes). This is contrasted to less than 10% if the value was below 2 log₁₀ *E. coli*/100 mL. It has been

suggested that monitoring of water quality in primary production for fecal indicator organisms such as *E. coli* can help inform farmers on deviations in good practices and situations that need corrective measures, thereby contributing to the assurance of a microbiologically safe product. This is especially the case since detection of pathogens in water (or fresh produce) is not always reliable for reasons described above, including statistical limitations, leading to a false sense of security.

The limit of an *E. coli* criterion in water (or fresh produce) is set according to what is generally obtainable when applying good practices and is not a direct indicator of risk. However, an increased number of *E. coli* cells (above the level normally observed) indicates a higher degree of exposure to fecal contamination from pathogen reservoirs and/or cross-contamination or growth (EFSA 2014). But indeed the utility of *E. coli* screening is debatable with regard to public health. For example, Ahmed and others (2010) found that 12% of roof harvest rainwater samples had <1 CFU *E. coli*/100 mL but were positive for one or more pathogens.

It has been suggested that the enterococci perform better than *E. coli* in terms of indicating fecal contamination and pathogen presence in environmental waters (perhaps because they are more environmentally persistent), although data are mixed (Kinzelman and others 2003; Lemarchand and Lebaron 2003; Hörman and others 2004; Harwood and others 2005). Alternative indicators such as *Bacteroides* spp., *Bifidobacterium* spp., and *Clostridium perfringens*, as well as bacteriophages (for example, coliphage MS2 or ΦX174 or *Bacteroides fragilis* phage B40-8) and adenoviruses, have also been proposed. Again, there are no compelling data about their utility to date.

The frequency of testing and the maximum allowed indicator level are still points of debate. Sampling once or twice a year provides some information on water quality, but eventually a high variability in the water quality, in particular for surface waters or during the growing season, may occur. Overall, the frequency of testing should depend upon the exact farm management and operation practices and climatic conditions. For instance, borehole water is less vulnerable to contamination and will demand less frequent testing than open reservoirs (of course, depending upon the construction of the reservoir). Furthermore, climate incidents such as flood, runoff of storm water, and so on would by necessity increase the frequency of testing. However, the presence of an effective and well-operated water treatment system implies a need for less frequent testing, which would be done merely for verification purposes.

Future Perspectives: the Role of Risk Assessment in Managing the Use of Water in Fresh Produce Primary Production

Guidelines and regulations dealing with microbial standards are often empirically designed, based upon prior experience on what is achievable under good practices, and has been shown to function by prior history and epidemiological evidence as appropriate in protecting consumers' health. But other strategies for managing health risks may also be effective (Carr and Blumenthal 2004). For example, the latest guidelines by the World Health Organization (2006) for use of wastewater in agriculture have been revised substantially by replacing the fecal coliform guideline with health-based targets defined through attributable risk and disability-adjusted life years (DALYs). As such, governments in developing countries have been given greater flexibility in achieving these targets (World Health Organization 2006). These guidelines are intended to be used as the basis for the development of na-

tional and international approaches to managing the health risk from hazards associated with treated wastewater use in agriculture. An example of implementation of this approach is illustrated in the AGWR report (O'Toole and others 2010). This study showed how to translate a health outcome target to performance targets for water treatment, and irrigation and farming practices. It shows how microbial risk assessment can be used in a regulatory framework to guide food producers to the appropriate risk management interventions (based on a combination of barriers) in the chain from irrigated fresh produce to consumer. Another example is a quantitative microbial risk assessment (QMRA) study elaborated in Sweden by Ottoson and others (2011). The QMRA indicated that reducing the maximum contamination level of irrigation water from 4 log₁₀ CFU to 2 log₁₀ CFU *E. coli*/100 mL would lead to a 5-fold reduction in verocytotoxin-producing *E. coli* illnesses due to consumption of iceberg lettuce. Besides controlling the microbiology of the irrigation water source, other recommendations could be made using this model, such as increasing the time between irrigation and harvest. Specifically, cessation of irrigation for, respectively, 1, 2, 4, and 7 days, which reduced the risk 3, 8.8, and 18 times. However, depending on the weather conditions, cessation of irrigation may not be possible in all cases.

Stine and others (2005) computed the maximum concentration of *Salmonella* and hepatitis A virus (HAV) in irrigation water that would result in a 10⁻⁴ annual risk of infection for individuals consuming different types of fresh produce that were irrigated under different conditions. Their findings indicated *Salmonella* concentrations could range from a low of 1.5 × 10² CFU/mL to a high of 7.2 × 10⁶ CFU/100 mL for furrow-irrigated lettuce, depending upon when the last irrigation event occurred (1 or 14 days before harvest, respectively). Hamilton and others (2006) developed a microbial risk assessment (MRA) model to estimate the risk of enteric virus illness when secondary effluent was used to irrigate horticultural crops (broccoli, cucumber, cabbage, and lettuce). The model computed the daily exposure based on the human body mass, daily consumption, virus concentration in water, volume of irrigation water deposited on the product, virus die-off, and time between last irrigation and harvest. A dose-response model for rotavirus was used as a proxy. Across the various produce crops, the annual risk of infection ranged from a low of 10⁻⁹ to 10⁻³ when irrigation using reclaimed water was ceased 2 weeks before harvest, to a high of 10⁻³ to 10⁻¹ when irrigation use was ceased 1 day before harvest.

Few site-specific data points were available for most of these MRAs, meaning that many assumptions were necessary. Specific parameters lacking hard data included the rates of pathogen transfer from irrigation water to crops, pathogen penetration in food crops, and pathogen survival on or in food crops. Data on these factors have been accumulating over the last decade, and this should improve the reliability of future MRA estimates. However, the sheer number of different fresh produce commodities and pathogens, combined with water sources and irrigation practices in different locations, means that developing risk models that can span the breadth of fresh produce safety will be a considerable challenge.

Overall Conclusions

Outbreaks of foodborne disease associated with fresh produce are not uncommon. The true disease burden is unknown due to under-reporting, the impact of sporadic disease, and poor epidemiological surveillance. There have been several outbreaks linked to contaminated irrigation water. Many different sources of water and methods are used for irrigation of fresh produce around the world.

There are 2 main sources of irrigation water: (i) surface water or treated wastewater (more prone to contamination and variable in water quality); and (ii) groundwater reserves or collected rainfall water (less prone to contamination or more controlled water quality if stored properly). Drip or subsurface irrigation limits direct contact between edible plant tissue and irrigation water (splashes) and thus is less likely to introduce pathogens than furrow or sprinkler irrigation. Codes of practice stress the importance of the quality of the irrigation water source for ensuring safety of fresh produce. A few general principles of preventive measures are (i) regular execution of and response to sanitary surveys; (ii) maintenance of irrigation water reservoirs and distribution systems; (iii) adequate water treatments to gain better water quality; and (iv) fecal indicator tests to monitor water quality. These measures are particularly helpful after climatic incidents. If working under conditions of good agricultural practices, in most cases, pathogen contamination of waters and fresh produce is expected to be an infrequent and temporary event, so direct pathogen screening of water (or produce) is likely to be ineffective. Nevertheless, this might be different in regions or under conditions in which contaminated surface water or insufficiently treated wastewater is used due to lack of access to clean water. Still, sanitary surveys and observational audits might also be more useful in these situations, as testing alone should never be relied upon as a food safety management tool, but rather should complement existing strategies (GAPs). An emerging alternative is the use of MRA to guide risk management directions for effective pathogen control and to select the most appropriate control measures to manage the use of water in fresh produce production.

Acknowledgments

This work was conducted by an expert group of the European branch of the International Life Sciences Institute (ILSI Europe). The authors would like to thank Dr Robert Bos (previously WHO, now IWA); Dr Sarah Cahill (FAO); Prof. Thor Axel Stenström (Swedish Institute for Infectious Disease Control); Dr Liqa Raschid-Sally (International Water Management Institute); Dr Annick Moreau and Dr Fabrice Peladan (Danone); Dr Michele Storrs Mabilat (bioMérieux); and these ILSI Branches (ILSI South Africa, ILSI Southeast Asia Region, ILSI India, ILSI Mexico, ILSI South Andean, ILSI North Andean, ILSI Korea, and ILSI Japan) for their active contributions to this study. The expert group received funding from the ILSI Europe Emerging Microbiological Issues Task Force. Industry members of this task force are listed on the ILSI Europe website at www.ils.eu. For further information about ILSI Europe, please email info@ilsieurope.be or call +32 2 771 00 14. The opinions expressed herein and the conclusions of this publication are those of the authors and do not necessarily represent the views of ILSI Europe nor those of its member companies.

References

- Ahmad I, Chua PC. 2013. Trends in production and trade of tropical fruits in ASEAN countries. *ISHS Acta Horticulturae* 975: IV International Symposium on Tropical and Subtropical Fruits. p 559–80.
- Ahmed W, Rodgers L, Sidhu JPS, Toze S. 2002. Fecal indicators and zoonotic pathogens in household drinking water taps fed from rainwater tanks in Southeast Queensland, Australia. *Appl Environ Microbiol* 78:219–26.
- Ahn JH, Grant SB, Surbeck CQ, DiGiacomo PM, Nezhlin NP, Jiang S. 2005. Coastal water quality impact of stormwater runoff from an urban watershed in southern California. *Environ Sci Technol* 39:5940–53.
- Amoah P, Drechsel P, Abaidoo RC, Klutse A. 2007. Effectiveness of common and improved sanitary washing methods in selected cities of West Africa for the reduction of coliform bacteria and helminth eggs on vegetables. *Trop Med Int Health* 12:40–50.
- Amoah P, Keraita B, Drechsel P, Abaidoo RC, Konradsen F, Akple M. 2011. Low-cost options for health risk reduction where crops are irrigated with polluted water in West Africa. Available from: http://www.iwmi.cgiar.org/Publications/IWMI_Research_Reports/PDF/PUB141/RR141.pdf. Accessed 2014 August 1.
- Arthurson V, Sessitsch A, Jäderlund L. 2011. Persistence and spread of *Salmonella enterica* serovar Weltevreden in soil and on spinach plants. *FEMS Microbiol Lett* 314:67–74.
- Asano T, Cotruvo JA. 2004. Groundwater recharge with reclaimed municipal wastewater: health and regulatory considerations. *Water Res* 38:1941–51.
- Astrom J, Pettersson T, Stenstrom T, Bergstedt O. 2009. Variability analysis of pathogen and indicator loads from urban sewer systems along a river. *Water Sci Technol* 59:203–12.
- Barker-Reid F, Harapas D, Engleitner S, Kreidl S, Holmes R, Faggian R. 2009. Persistence of *Escherichia coli* on injured iceberg lettuce in the field, overhead irrigated with contaminated water. *J Food Prot* 72:458–64.
- Barno A, Ondanje B, Ngwiri J. 2009. Dynamics of horticultural export to European Union market: challenges and opportunities in Sub-Saharan Africa. *ISHS Acta Horticulturae* 911: I All Africa Horticultural Congress. p 61–72.
- Bashour I, Nimah M. 2004. Fertigation potentials in the Near East region. IPI Regional Workshop Proceedings on Potassium and Fertigation Development in West Asia and North Africa; Rabat, Morocco, 2004 November 24–28.
- Basu P, Scholten BA. 2012. Crop livestock systems in rural development: linking India's Green and White Revolutions. *Intl J Agr Sustain* 10:175–91.
- Beatty ME, LaPorte TN, Phan Q, Van Duyn SV, Braden C. 2004. A multistate outbreak of *Salmonella enterica* serotype Saintpaul infections linked to mango consumption: a recurrent theme. *Clin Infect Dis* 38:1337–8.
- Behravesh B, Mody RK, Jungk J, Gaul L, Redd JT, Chen S, Cosgrove S, Hedican E, Sweat D, Chávez-Hausler L, Snow SL, Hanson, H, Nguyen TA, Sodha SV, Boore AL, Russo E, Mikoleit M, Theobald L, Gerner-Smidt P, Hoekstra RM, Angulo FJ, Swerdlow DL, Tauxe RV, Griffin PM, Williams IT. 2011. 2008 Outbreak of *Salmonella* Saintpaul infections associated with raw produce. *N Engl J Med* 364:918–27.
- Betts R. 2014. Microbial update: fruit and salad. *Intl Food Hyg* 25:9–12.
- Beuchat LR. 2002. Ecological factors influencing survival and growth of human pathogens on raw fruits and vegetables. *Microbes Infect* 4:413–23.
- Blumenthal UJ, Mara DD, Peasley A, Ruiz-Palacios G, Stott R. 2000. Guidelines for the microbiological quality of treated wastewater used in agriculture: recommendations for revising WHO guidelines. *Bull World Health Organ* 78:1104–16.
- BOE. 2007. Spanish Royal Decision, 1620/2007: Royal Decree where is regulated the legal regime for reuse of purified waste waters. *Boletín Oficial del Estado* No. 294.
- Brackett RE. 1999. Incidence, contributing factors, and control of bacterial pathogens in produce. *Postharvest Biol Technol* 15:305–11.
- Brandl MT, Amundson R. 2008. Leaf age as a risk factor in contamination of lettuce with *Escherichia coli* O157:H7 and *Salmonella enterica*. *Appl Environ Microbiol* 74:2298–306.
- Brandl MT. 2006. Fitness of human enteric pathogens on plants and implications for food safety 1. *Annu Rev Phytopathol* 44:367–92.
- Brandl MT, Mandrell RE. 2002. Fitness of *Salmonella enterica* serovar Thompson in the cilantro phyllosphere. *Appl Environ Microbiol* 68:3614–21.
- Burch JD, Thomas KE. 1998. Water disinfection for developing countries and potential for solar thermal pasteurization. *Solar Energy* 64:87–97.
- Cardenas C, Molina K, Heredia N, García S. 2013. Evaluation of microbial contamination of tomatoes and peppers at retail markets in Monterrey, Mexico. *J Food Prot* 76:1475–9.
- Carr R, Blumenthal U. 2004. Guidelines for the safe use of wastewater in agriculture: revisiting WHO guidelines. *Water Sci Technol* 50:31–8.
- Castro-Rosas J, Cerna-Cortés JF, Méndez-Reyes E, Lopez-Hernandez D, Gómez-Aldapa CA, Estrada-García T. 2012. Presence of faecal coliforms, *Escherichia coli* and diarrheagenic *E. coli* pathotypes in ready-to-eat salads, from an area where crops are irrigated with untreated sewage water. *Intl J Food Microbiol* 156:176–80.

- Centers for Disease Control and Prevention. 1997. Outbreaks of *Escherichia coli* O157:H7 infection and cryptosporidiosis associated with drinking unpasteurized apple cider. *MMWR Morb Mortal Wkly Rep* 46: 4–8.
- Centers for Disease Control and Prevention. 2011. Investigation update: multistate outbreak of listeriosis linked to whole cantaloupes from Jensen Farms. Available from: http://www.cdc.gov/listeria/outbreaks/cantaloupes-jensen-farms/index.html?cid=cs_654. Accessed 2014 September 25.
- Centers for Disease Control and Prevention. 2013. Investigation update: multistate outbreak of cyclosporiasis. Update. Available from: <http://www.cdc.gov/parasites/cyclosporiasis/outbreaks/investigation-2013.html>. Accessed 2014 September 25.
- Cevallos-Cevallos JM, Danyluk MD, Gu G, Vallad GE, Van Bruggen AHC. 2012. Dispersal of *Salmonella* Typhimurium by rain splash onto tomato plants. *J Food Prot* 75:472–9.
- CFERT, California Department of Health Services and U.S. Food And Drug Administration. 2007. Investigation of an *Escherichia coli* O157:H7 outbreak associated with Dole pre-packaged spinach. Available from: http://www.cdc.gov/nced/ehs/Docs/Investigation_of_an_E_Coli_Outbreak_Associated_with_Dole_Pre-Packaged_Spinach.pdf. Accessed 2014 September 25.
- Cirelli, GL, Consoli S, Di Grande V. 2008. Long-term storage of reclaimed water: the case studies in Sicily (Italy). *Desalination* 218: 62–73.
- Codex Alimentarius Commission. 2003b. Code of hygienic practice for fresh fruits and vegetables. Available from: <http://www.justice.gov.md/upload/Baza%20de%20date/Materiale%202009/Legislatie/Codex%20Stan%2053-2003.pdf>. Accessed 2014 September 25.
- Codex Alimentarius Commission. 2003c. Recommended international code of practice: General principles of food hygiene. Recommended International Code of Practice General Principles of Food Hygiene. Available from: <http://www.mhlw.go.jp/english/topics/importedfoods/guideline/dl/04.pdf>. Accessed 2014 September 25.
- Codex Alimentarius Commission. 2003a. Code of hygienic practice for fresh fruits and vegetables. CAC/RCP 53-2003. Available from: ftp://ftp.fao.org/codex/publications/Booklets/FreshFruitsVeg/FFV_2007_EN.pdf. Accessed 2014 September 25.
- Codex Alimentarius Commission. 2012. Guidelines on the Application of General Principles of Food Hygiene to the Control of Viruses in Food. Available from: http://www.codexalimentarius.org/input/download/standards/13215/CXG_079e.pdf. Accessed 2014 September 25.
- Cooley MB, Chao D, Mandrell RE. 2006. *Escherichia coli* O157:H7 survival and growth on lettuce is altered by the presence of epiphytic bacteria. *J Food Prot* 69:2329–35.
- Dauchet L, Czernichow S, Bertrais S, Blacher J, Galan P, Hercberg S. 2005. Fruits and vegetables intake in the SU.VI.MAX study and systolic blood pressure change. *Arch Mal Coeur Vaiss* 99:669–73.
- de Quadros Rodrigues R, Loikoa MR, Minéia Daniel de Paula C, Hessela CT, Jacxsens L, Uyttendaele M, Renar JB, Tondo EC. 2014. Microbiological contamination linked to implementation of good agricultural practices in the production of organic lettuce in Southern Brazil. *Food Control* 42:152–164.
- De Roeber C. 1998. Microbiological safety evaluations and recommendations on fresh produce. *Food Control* 9:321–47.
- Delaquis P, Bach S, Dinu LD. 2007. Behavior of *Escherichia coli* O157:H7 in leafy vegetables. *J Food Prot* 70:1966–74.
- Dell'Erba, A, Falsanisi D, Liberti L, Notarnicola M, Santoro D. 2004. Disinfecting behaviour of peracetic acid for municipal wastewater reuse. *Desalination* 168:435–42.
- DeWaal CS, Tian XA, Plunkett D. 2009. Outbreak alert! Analyzing foodborne outbreaks 1998–2007. *Outbreak Alert!* Available from: www.cspinet.org/new/pdf/outbreakalertreport09.pdf. Accessed 2014 September 25.
- Döller PC, Dietrich K, Philipp N, Brockmann S, Dreweck C, Vonthein R, Wagner-Wiening C, Weidenmann R. 2000. Cyclosporiasis outbreak in Germany associated with the consumption of salad. *Emerg Infect Dis* 8:992–4.
- E.P.H.C. 2006. Australian guidelines for water recycling. Managing health and environmental risks. Phase 1. National water quality management strategy 21. Natural Resource Management Ministerial Council. Environment Protection and Heritage Council, Australian Health Ministers Conference Council, Canberra, Australia. Available from: <http://www.environment.gov.au/resource/national-water-quality-management-strategy-australian-guidelines-water-recycling-managing-0>. Accessed 2014 September 25.
- Edelstein M, Sundborger C, Hergens MP, Ivarsson S, Dryselius R, Insulander M, Jernberg C, Hutin Y, Wallensten A. 2013. Barriers to trace-back in a salad-associated EHEC outbreak, Sweden, June 2013. *PLoS currents* 6. Available from: <http://currents.plos.org/outbreaks/article/barriers-to-trace-back-in-a-salad-associated-ehec-outbreak-sweden-june-2013/>. Accessed 2014 September 25.
- EFSA. 2012. The European Union summary report on trends and sources of zoonoses, zoonotic agents and food-borne outbreaks in 2010. *EFSA J* 10:1–442.
- EFSA. 2014. The community summary report on trends and sources of zoonoses, zoonotic agents and food-borne outbreaks in the European Union in 2011. *EFSA J* 12:3547.
- EFSA Panel on Biological Hazards (BIOHAZ). 2013. Scientific opinion on VTEC-serotype and scientific criteria regarding pathogenicity assessment. *EFSA J* 11:1–106.
- EFSA Panel on Biological Hazards (BIOHAZ) Panel. 2012. Scientific opinion on the risk posed by pathogens in food of non-animal origin. Part 1 (outbreak data analysis and risk ranking of food/pathogen combinations). *EFSA J* 11:1–138.
- Enriquez C, Alum A, Suarez-Rey EM, Choi CY, Oron G, Gerba CP. 2003. Bacteriophages MS2 and PRD1 in turfgrass by subsurface drip irrigation. *J Environ Eng* 129:852–7.
- Erickson MC, Webb CC, az-Perez JC, Phatak SC, Silvoy JJ, Davey L, Payton AS, Liao J, Ma L, Doyle MP. 2010. Surface and internalized *Escherichia coli* O157:H7 on field-grown spinach and lettuce treated with spray-contaminated irrigation water. *J Food Prot* 73:1023–9.
- Ethelberg S, Lisby M, Bottiger B, Schultz AC, Villif A, Jensen T, Olsen KE, Scheutz F, Kjelson C, Muller L. 2010. Outbreaks of gastroenteritis linked to lettuce, Denmark, January 2010. *Eurosurveillance* 15.
- European Commission. 2004. Regulation (EC) No 852/2004 of the European Parliament and of the Council of 29 April 2004. Available from: <http://eur-lex.europa.eu/LexUriServ/LexUriServ.do?uri=OJ:L:2004:139:0001:0054:en:PDF>. Accessed 2014 September 25.
- Eurostat. 2013. Agri-environmental indicator – irrigation – Statistics Explained (2013/3/5) HYPERLINK “http://epp.eurostat.ec.europa.eu/statistics_explained/index.php/Agri-environmental_indicator_-_irrigation%20accessed%20on%2011/12/2013” http://epp.eurostat.ec.europa.eu/statistics_explained/index.php/Agri-environmental_indicator_-_irrigation accessed on 11/12/2013.
- Evans CA, Coombes PJ, Dunstan RH. 2006. Wind, rain and bacteria: the effect of weather on the microbial composition of roof-harvested rainwater. *Water Res* 40:37–44.
- Falkenhorst G, Krusell L, Lisby M, Madsen SB, Böttiger B, Mølbak K. 2005. Imported frozen raspberries cause a series of norovirus outbreaks in Denmark, 2005. *Eurosurveillance* 10.
- Falsanisi D, Gehr R, Santoro D, Dell'Erba A, Notarnicola M, Liberti L. 2006. Kinetics of PAA demand and its implications on disinfection of wastewaters. *Wat Qual Res J Can* 41:398–409.
- Fattal B, Margalith M, Shuval HI, Wax Y, Morag A. 1987. Viral antibodies in agricultural populations exposed to aerosols from wastewater irrigation during a viral disease outbreak. *Am J Epidemiol* 125:899–906.
- FDA. 1998. Guide to minimize microbial food safety hazards for fresh fruits and vegetables. Available from: <http://www.fda.gov/downloads/Food/GuidanceRegulation/UCM169112.pdf>. Accessed 2014 September 25.
- FDA. 2009a. Draft guidance for industry: guide to minimize microbial food safety hazards of leafy greens. Available from: <http://www.fda.gov/food/guidanceregulation/guidancedocumentsregulatoryinformation/produceplantproducts/ucm174200.htm>. Accessed 2014 September 25.
- FDA. 2009b. Draft guidance for industry: guide to minimize microbial food safety hazards of melons. Available from: <http://www.fda.gov/Food/GuidanceRegulation/GuidanceDocumentsRegulatoryInformation/ProducePlantProducts/ucm174171.htm>. Accessed 2014 September 25.
- FDA. 2009c. Draft guidance for industry: guide to minimize microbial food safety hazards of tomatoes. Available from: <http://www.fda.gov/Food/GuidanceRegulation/GuidanceDocumentsRegulatoryInformation/ProducePlantProducts/ucm173902.htm>. Accessed 2014 September 25.
- FDA. 2009d. Draft guidance for industry: guide to minimize microbial food safety hazards of leafy greens. Available from: <http://www.fda.gov/food/guidanceregulation/guidancedocumentsregulatoryinformation/produceplantproducts/ucm174200.htm>. Accessed 2014 September 25.
- Feenstra S, Hussain R, Van der Hoek W. 2000. Health risks of irrigation with untreated urban wastewater in the southern Punjab, Pakistan. Available

- from: <http://publications.iwmi.org/pdf/H026997.pdf>. Accessed 2014 September 25.
- Fett WF 2000. Naturally occurring biofilms on alfalfa and other types of sprouts. *J Food Prot* 63:625–32.
- Fong TT, Mansfield LS, Wilson DL, Schwab DJ, Molloy SL, Rose JB. 2007. Massive microbiological groundwater contamination associated with a waterborne outbreak in Lake Erie, South Bass Island, Ohio. *Environ Health Perspect* 115:856–64.
- Fonseca JM, Fallon SD, Sanchez CA, Nolte KD. 2011. *Escherichia coli* survival in lettuce fields following its introduction through different irrigation systems. *J Appl Microbiol* 110:893–902.
- Foster, T., Brozovic, N., Butler, A.P. 2014. Modeling irrigation behavior in groundwater systems. *Water resources research* 50: 6370–6389.
- Franz E, Semenov AV, Termorshuizen AJ, De Vos OJ, Bokhorst JG, van Bruggen AHC. 2008. Manure-amended soil characteristics affecting the survival of *E. coli* O157:H7 in 36 Dutch soils. *Environ Microbiol* 10: 313–27.
- Franz E, van Diepeningen AD, de Vos OJ, van Bruggen AHC. 2005. Effects of cattle feeding regimen and soil management type on the fate of *Escherichia coli* O157:H7 and *Salmonella enterica* serovar Typhimurium in manure, manure-amended soil, and lettuce. *Appl Environ Microbiol* 71:6165–74.
- Geldreich EE. 1991. Microbial water quality concerns for water supply use. *Environ Toxicol Wat Qual* 6:209–23.
- Gerba CP, Smith EE. 2005. Sources of pathogenic microorganisms and their fate during land application of wastes. *J Environ Qual* 34:42–8.
- Gerba CP, Choi C. 2006. Role of irrigation water in crop contamination by viruses. In: Goyal S M, editor. *Virusus in foods*. New York: Springer. p 257–63.
- Ghadiri H, Dordipour I, Bybordi M, Malakouti MJ. 2006. Potential use of Caspian Sea water for supplementary irrigation in Northern Iran. *Agr Wat Manag* 79:209–24.
- Ghassemi F, Jakeman AJ, Nix HA. 1995. Salinisation of land and water resources: human causes, extent, management and case studies. Wallingford: CAB International.
- Gibbs R, Pingault N, Mazzucchelli T, O'Reilly L, MacKenzie B, Green J, Mogyrosy R, Stafford R, Bell R, Hiley L. 2009. An outbreak of *Salmonella enterica* serotype Litchfield infection in Australia linked to consumption of contaminated papaya. *J Food Prot* 72:1094–8.
- Gil MI, Selma MV, Suslow T, Jacxsens L, Uyttendaele M, Allende A. 2015. Pre- and post-harvest preventive measures and intervention strategies to control microbial food safety hazards of fresh leafy vegetables. *Crit Rev Food Sci Nutr* 55:453–468.
- Girardin H, Morris CE, Albagnac C, Dreux N, Glaux C. 2005. Behaviour of the pathogen surrogates *Listeria innocua* and *Clostridium sporogenes* during production of parsley in fields fertilized with contaminated amendments. *FEMS Microbiol Ecol* 54:287–95.
- Gleick PH. 2000. The world's water 2000–2001: the biennial report on freshwater resources. Washington, DC: Island Press.
- Global Gap. 2013. Integrated farm assurance/all farm base/crops base/fruit and vegetables: control points and compliance criteria. Available from: http://www.globalgap.org/export/sites/default/.content/.galleries/.documents/130315_gg_ifa_cpcc_af_cb_fw_v4_0--2_en.pdf. Accessed 2014 September 25.
- González M, Hänninen ML. 2012. Effect of temperature and antimicrobial resistance on survival of *Campylobacter jejuni* in well water: application of the Weibull model. *J Appl Microbiol* 113:84–293.
- Gorton M, Sauer J, Supatpongkul P. 2011. Wet markets, supermarkets and the “Big Middle” for food retailing in developing countries: evidence from Thailand. *World Develop* 39:1624–37.
- Goyal, SM, Gerba CP, Melnick JL. 1977. Occurrence and distribution of bacterial indicators and pathogens in canal communities along the Texas coast. *Appl Environ Microbiol* 34:139–49.
- Gras MH, Druet MC, Cerf O. 1994. Bacterial flora of salad leaves [ready-to-eat raw vegetables, biofilm, decontamination, scarole, frisee, chioGGia]. *Sci Aliment* 14: 173–88.
- Greene SK, Daly ER., Talbot EA, Demma LJ, Holzbauer S, Patel NJ, Hill TA, Walderhaug MO, Hoekstra RM, Lynch MF. 2008. Recurrent multistate outbreak of *Salmonella* Newport associated with tomatoes from contaminated fields, 2005. *Epidemiol Infect* 136:157–65.
- Guler E, Ozakdag D, Arda M, Yuksel M, Kabay N. 2010. Effect of temperature on seawater desalination–water quality analyses for desalinated seawater for its use as drinking and irrigation water. *Environ Geochem Health* 32:335–9.
- Hallam NB, West JR, Forster CF, Simms J. 2001. The potential for biofilm growth in water distribution systems. *Water Res* 35:4063–71.
- Hamilton AJ, Stagnitti F, Premier R, Boland AM, Hale G. 2006. Quantitative microbial risk assessment models for consumption of raw vegetables irrigated with reclaimed water. *Appl Environ Microbiol* 72:3284–90.
- Hanning IB, Nutt JD, Ricke SC. 2009. Salmonellosis outbreaks in the United States due to fresh produce: sources and potential intervention measures. *Foodborne Pathog Dis* 6:635–48.
- Harwood VJ, Levine AD, Scott TM, Chivukula V, Lukasik J, Farrah SR, Rose JB. 2005. Validity of the indicator organism paradigm for pathogen reduction in reclaimed water and public health protection. *Appl Environ Microbiol* 71:3163–70.
- Heaton JC, Jones K. 2008. Microbial contamination of fruit and vegetables and the behaviour of enteropathogens in the phyllosphere: a review. *J Appl Microbiol* 104:613–26.
- Herwaldt BL, Ackers ML. 1997. An outbreak in 1996 of cyclosporiasis associated with imported raspberries. *N Engl J Med* 336:1548–56.
- Hoang LMN, Fyfe M, Ong C, Harb J, Champagne S, Dixon B, Isaac-Renton J. 2005. Outbreak of cyclosporiasis in British Columbia associated with imported Thai basil. *Epidemiol Infect* 133:23–7.
- Holvoet K, De Keuckelaere A, Sampers I, Van Haute S, Stals A, Uyttendaele M. 2014. Quantitative study of cross-contamination with *Escherichia coli*, *E. coli* O157, MS2 phage and murine norovirus in a simulated fresh-cut lettuce wash process. *Food Control* 37:218–27.
- Horman A, Rimhanen-Finne R, Maunula L, von Bonsdorff CH, Torvela N, Heikinheimo A, Hanninen ML. 2004. *Campylobacter* spp., *Giardia* spp., *Cryptosporidium*, noroviruses and indicator organisms in surface water in southwestern Finland, 2000–2001. *Appl Environ Microbiol* 70: 87–95.
- Hunt RJ, Coplen TB, Haas NL, Saad DA, Borchardt MA. 2005. Investigating surface water–well interaction using stable isotope ratios of water. *J Hydrol* 302:154–72.
- Huong PTT, Everaarts AP, Neeteson JJ, Struik PC. 2013. Vegetable production in the Red River delta of Vietnam. I. Opportunities and constraints. *NJAS-Wageningen J Life Sci* 67:27–36.
- Hussain I., L. Raschid, M. A. Hanjra, F. Marikar, W. van der Hoek. 2002. Wastewater use in agriculture: Review of impacts and methodological issues in valuing impacts. Working Paper 37. Colombo, Sri Lanka: International Water Management Institute. ISBN 92-9090-472-0.
- Hutchison ML, Avery SM, Monaghan JM. 2008. The air-borne distribution of zoonotic agents from livestock waste spreading and microbiological risk to fresh produce from contaminated irrigation sources. *J Appl Microbiol* 105:848–57.
- Hutchison ML, Walters LD, Moore T, Thomas DJ, Avery SM. 2005. Fate of pathogens present in livestock wastes spread onto fescue plots. *Appl Environ Microbiol* 71:691–6.
- Ibekwe AM, Shouse PJ, Grieve CM. 2006. Quantification of survival of *Escherichia coli* O157:H7 on plants affected by contaminated irrigation water. *Eng Life Sci* 6:566–72.
- Ibenyassine K, Mhand RA, Karamoko Y, Anajjar B, Chouibani MM, Ennaji M. 2007. Bacterial pathogens recovered from vegetables irrigated by wastewater in Morocco. *J Environ Health* 69:47–51.
- Insulander M, Svenungsson B, Lebbad M, Karlsson L, de Jong B. 2010. A foodborne outbreak of *Cyclospora* infection in Stockholm, Sweden. *Foodborne Pathog Dis* 7:1585–7.
- IPTRID. 2001. Field guide on irrigated agriculture for field assistants. Available from: <ftp://ftp.fao.org/agl/iptrid/FieldGuideSourceFig.pdf>. Accessed 2014 September 25.
- Islam M, Doyle MP, Phatak SC, Millner P, Jiang X. 2004. Persistence of enterohemorrhagic *Escherichia coli* O157:H7 in soil and on leaf lettuce and parsley grown in fields treated with contaminated manure composts or irrigation water. *J Food Prot* 67:1365–70.
- Jacxsens L. 2010. Autocontrolegids aardappelen, groenten en fruit verwerkende industrie en handel. Versie 2 (in samenwerking met Belgapom, Vegebe en Fresh Trade Belgium).
- James J. 2006. Overview of Microbial Hazards in Fresh Fruit and Vegetables Operations, in *Microbial Hazard Identification in Fresh Fruit and Vegetables*. John Wiley & Sons, Inc. Hoboken. NJ. USA.
- Janisiewicz WJ, Conway WS, Brown, MW, Sapers GM, Fratamico P, Buchanan RL. 1999. Fate of *Escherichia coli* O157:H7 on fresh-cut apple tissue and its potential for transmission by fruit flies. *Appl Environ Microbiol* 65:1–5.

- Jiang HXL, Dian H, Chen ZL, Zhang L, Wang, XM, Chen JR, Liu YH, Liao XP, Liu JH, Zeng ZL. 2011. High prevalence and widespread distribution of multi-resistant *Escherichia coli* isolates in pigs and poultry in China. *Veterinary J* 187:99–103.
- Johannessen GS, Bengtsson GB, Heier BT, Bredholt S, Wasteson Y, Rørvik LM. 2005. Potential uptake of *Escherichia coli* O157:H7 from organic manure into crisphead lettuce. *Appl Environ Microbiol* 71:2221–5.
- Johnston LM, Jaykus LA, Moll D, Anciso J, Mora B, Moe CL. 2006. A field study of the microbiological quality of fresh produce of domestic and Mexican origin. *Intl J Food Microbiol* 112:83–95.
- Keraita B, Drechsel P, Huibers F, Raschid-Sally L. 2002. Wastewater use in informal irrigation in urban and peri-urban areas of Kumasi, Ghana. *Urban Agr Mag* 8:11–3.
- Keraita B, Drechsel P, Konradsen F. 2010. Up and down the sanitation ladder: harmonizing the treatment and multiple-barrier perspectives on risk reduction in wastewater irrigated agriculture. *Irrig Drain Sys* 24:23–35.
- Keraita B, Konradsen F, Drechsel P, Abaidoo, RC. 2007. Reducing microbial contamination on wastewater-irrigated lettuce by cessation of irrigation before harvesting. *Trop Med Intl Health* 12:8–14.
- Khalil R, Gomaa M. 2014. Evaluation of the microbiological quality of conventional and organic leafy greens at the time of purchase from retail markets in Alexandria, Egypt. *Pol J Microbiol* 63:237–43.
- Kinzelman, J., Ng, C., Jackson, E., Gradus, S. and Bagley, R. (2003). Enterococci as indicators of Lake Michigan recreational water quality: comparison of two methodologies and their impacts on public health regulatory events. *Appl Environ Microbiol* 69: 92–96.
- Kisluk G, Yaron S. 2012. Presence and persistence of *Salmonella enterica* serotype Typhimurium in the phyllosphere and rhizosphere of spray-irrigated parsley. *Appl Environ Microbiol* 78:4030–6.
- Koivunen J, Heinonen-Tanski H. 2005. Peracetic acid (PAA) disinfection of primary, secondary and tertiary treated municipal wastewaters. *Water Res* 39:4445–53.
- Krinner W, Lallana C, Estrela T, Nixon S, Zabel T, Laffon L, Rees G, Cole G. 1999. Sustainable water use in Europe: Part 1. Sectoral use of water, Environmental assessment report. http://www.eea.europa.eu/publications/binarveenviasses01pdf/at_download/file. Accessed 2014 August 14.
- Leifert C, Ball K, Volakakis N, Cooper JM. 2008. Control of enteric pathogens in ready-to-eat vegetable crops in organic and low input production systems: a HACCP-based approach. *J Appl Microbiol* 105:931–50.
- Lemarchand K, Lebaron P. 2003. Occurrence of salmonella spp and cryptosporidium spp in a French coastal watershed: relationship with fecal indicators. *FEMS Microbiological Letters* 218:203–9.
- Leon JS, Jaykus LA, Moe CL. 2009. Microbiology of fruits and vegetables. Microbiologically safe foods. John Wiley & Sons, Inc., Hoboken, NJ. USA. p 255–90.
- LGMA, Leafy Greens Marketing Agreement. 2012. Commodity specific food safety guidelines for the production and harvest of lettuce and leafy greens. Available from: <http://www.caleafygreens.ca.gov/sites/default/files/01.20.12%20CALGMA%20GAP%20-%20metrics.pdf>. Accessed 2014 September 25.
- Liberti L, Notarnicola M, Lopez A. 2000. Advanced treatment for municipal wastewater reuse in agriculture: III. Ozone disinfection. *Ozone Sci Eng* 22:151–66.
- Liberti L, Notarnicola M, Boghetich G, Lopez, A. 2001. Advanced treatment for municipal wastewater reuse in agriculture. UV disinfection: bacteria inactivation. *Aqua* 50:275–85.
- Liu C, Hofstra N, Franz E. 2013. Impacts of climate change on the microbial safety of pre-harvest leafy green vegetables as indicated by *Escherichia coli* O157 and *Salmonella* spp. *Intl J Food Microbiol* 163:119–28.
- Lucena F, Ribas F, Duran AE, Skraber S, Gantzer C, Campos C, Moron A, Calderon E, Jofre J. 2006. Occurrence of bacterial indicators and bacteriophages infecting enteric bacteria in groundwater in different geographical areas. *J Appl Microbiol* 101:96–102.
- Lynch MF, Tauxe RV, Hedberg CW. 2009. The growing burden of foodborne outbreaks due to contaminated fresh produce: risks and opportunities. *Epidemiol Infect* 137:307–15.
- Marites M, Tiengco N, Bidwell K. 2010. Risk analysis integrating livelihood and economic impacts of wastewater irrigation on health. Wastewater irrigation and health: assessing and mitigating risk in low-income countries. London: Earthscan-International Development Research Centre. p 127–48.
- Matmerk KSL. 2010. The Norwegian Agricultural Quality System and Food Branding Foundation. Guideline 5. Vegetables, fruits, soft fruits and greenhouses. Available from: [http://www.matmerk.no/kunder/matmerk/mm.nsf/lupgraphics/Rettleiar%205%20Gr%C3%B8nsaker,%20frukt,%20b%20C3%A6r,%20planteskule,%20blomstrar%20og%20vekstus%20-%20nyno.%20-%202014.pdf](http://www.matmerk.no/kunder/matmerk/mm.nsf/lupgraphics/Rettleiar%205%20Gr%C3%B8nsaker,%20frukt,%20b%20C3%A6r,%20planteskule,%20blomstrar%20og%20vekstus%20-%20nyno.%20-%202014.pdf/$file/Rettleiar%205%20Gr%C3%B8nsaker,%20frukt,%20b%20C3%A6r,%20planteskule,%20blomstrar%20og%20vekstus%20-%20nyno.%20-%202014.pdf). Accessed 2014 October 30.
- McDonald's Good Agricultural Practices. Food Safety Standards. 2011. Issued March 14, 2011 Version 8.3. By McDonald's Worldwide Quality Systems, 60 pages
- Mdluli F, Thamaga-Chitja J, Schmidt S. 2013. Appraisal of hygiene indicators and farming practices in the production of leafy vegetables by organic small-scale farmers in uMbumbulu (Rural KwaZulu-Natal, South Africa). *Intl J Environ Res Public Health* 10:4323–38.
- Medina MS, Tudela JA, Marín A, Allende A, Gil MI. 2012. Short postharvest storage under low relative humidity improves quality and shelf life of minimally processed baby spinach (*Spinacia oleracea* L.). *Postharvest Biol Technol* 67:1–9.
- Melloul AA, Hassani L, Rafouk L. 2001. Salmonella contamination of vegetables irrigated with untreated wastewater. *World J Microbiol Biotechnol* 17:207–9.
- Mena KD. 2006. Produce quality and foodborne disease: assessing water's role. Microbial hazard identification in fresh fruit and vegetables. John Wiley & Sons, Inc., Hoboken, NJ. USA. p 95–114.
- Minhas PS, Sharma N, Yadav RK, Joshi PK. 2006. Prevalence and control of pathogenic contamination in some sewage irrigated vegetable, forage and cereal grain crops. *Bioresour Technol* 97:1174–8.
- Mootian G, Wu WH, Matthews KR. 2009. Transfer of *Escherichia coli* O157:H7 from soil, water, and manure contaminated with low numbers of the pathogen to lettuce plants. *J Food Prot* 72:2308–12.
- Morris CE, Monier JM. 2003. The ecological significance of biofilm formation by plant-associated bacteria. *Annu Rev Phytopathol* 41:429–53.
- Mossel DAA. 1978. Index and indicator organisms—a current assessment of their usefulness and significance [food microbiology]. *Food Technol Aust* 30:212–9.
- Mossel DAA. 1983. Marker (index and indicator) organisms in food and drinking water. Semantics, ecology, taxonomy and enumeration. *Antonie van Leeuwenhoek* 48:609–11.
- Moyne AL, Sudarshana MR, Blessington T, Koike ST, Cahn MD, Harris LJ. 2011. Fate of *Escherichia coli* O157:H7 in field-inoculated lettuce. *Food Microbiol* 28:1417–25.
- Muñoz I, Fernández-Alba AR. 2008. Reducing the environmental impacts of reverse osmosis desalination by using brackish groundwater resources. *Water Res* 42:801–11.
- Ndiaye ML, Niang S, Pfeifer H, Peduzzi R, Tonolla M, Dieng Y. 2011. Effect of irrigation water and processing on the microbial quality of lettuces produced and sold on markets in Dakar (Senegal). *IrrigDrain* 60:509–17.
- New England Farmers Union. 2013. Food Safety Modernization Act: Proposed produce safety rules: Microbial standards for water (Agricultural Water Requirements). Available from: <http://www.regulations.gov/#!documentDetail;D=FDA-2011-N-0921-0376>. Accessed 2014 August 14.
- Ntahimpera N, Wilson LL, Ellis MA, Madden LV. 1999. Comparison of rain effects on splash dispersal of three *Colletotrichum* species infecting strawberry. *Phytopathology* 89:555–63.
- Nygård K, Lassen J, Vold L, Andersson Y, Fisher I, Löfdahl S, Threlfall J, Luzzi I, Peters T, Hampton M. 2008. Outbreak of *Salmonella* Thompson infections linked to imported rucola lettuce. *Foodborne Pathog Dis* 5:165–73.
- O'Toole J, Sinclair M, Leder K. 2010. Quantitative microbial risk assessment and Australian Guidelines for Water Recycling: two case studies. *Food Aust* 62:408–412.
- Obuobie E, Keraita B, Danso G, Amoah P, Cofie OO, Raschid-Sally L, Drechsel P. 2010. Irrigated urban vegetable production in Ghana: Characteristics, benefits and risks. Available from: http://www.iwmi.cgiar.org/Publications/Books/PDF/irrigated_urban_vegetable_production_in_ghana.pdf. Accessed 2014 August 14.
- Olaimat AN, Holley RA. 2012. Factors influencing the microbial safety of fresh produce: a review. *Food Microbiol* 32:1–19.
- Oliveira M, Usall J, Viñas I, Solsona C, Abadías M. 2011. Transfer of *Listeria innocua* from contaminated compost and irrigation water to lettuce leaves. *Food Microbiol* 28:590–6.
- Oliveira M, Viñas I, Usall J, Anguera M, Abadías M. 2012. Presence and survival of *Escherichia coli* O157:H7 on lettuce leaves and in soil treated with contaminated compost and irrigation water. *Intl J Food Microbiol* 156:133–140.
- Ongley E. 1996. Control of water pollution from agriculture—FAO irrigation and drainage paper 55. Available from: <http://www.fao.org/3/a-w2598e/index.html>. Accessed 2014 August 14.

- Oron G, Armon R, Mandelbaum R, Manor Y, Campos C, Gillerman L, Salgot M, Gerba C, Klein I, Enriquez C. 2001. Secondary wastewater disposal for crop irrigation with minimal risks. *Water Sci Technol* 43:139–46.
- Ottoson JR, Nyberg K, Lindqvist R, Albiñ A. 2011. Quantitative microbial risk assessment for *Escherichia coli* O157 on lettuce, based on survival data from controlled studies in a climate chamber. *J Food Prot* 74: 2000–7.
- Parker JK, McIntyre D, Noble RT. 2010. Characterizing fecal contamination in stormwater runoff in coastal North Carolina, USA. *Water Res* 44:4186–94.
- Pedrero F, Kalavrouziotis I, Alarcón JJ, Koukoulakis P, Asano T. 2010. Use of treated municipal wastewater in irrigated agriculture. Review of some practices in Spain and Greece. *Agr Water Manag* 97:1233–41.
- Pereira LS, Oweis T, Zairi A. 2002. Irrigation management under water scarcity. *Agr Water Manag* 57:175–206.
- Pietravalle S, Van den Bosch F, Welham SJ, Parker SR, Lovell DJ. 2001. Modelling of rain splash trajectories and prediction of rain splash height. *Agr Forest Meteorol* 109:171–85.
- Pillai SD. 1998. Microbial pathogens within aquifers: principles and protocols. Michigan: University of Michigan Press.
- Primary Production. 2013. Module A: Primaire plantaardige productie. Versie 1.1. Available from: http://www.primaryproduction.be/fileadmin/files/G-040_Module_A_NL.131030.pdf. Accessed 2014 August 14.
- Public Health England. 2014. VTEC outbreak(s) linked to watercress: an update. Available from: https://www.gov.uk/government/uploads/system/uploads/attachment_data/file/326918/hpr0714.pdf. Accessed 2014 August 14.
- Qadir M. 2008. Sustainable Management of Wastewater for Agriculture: Proceedings of the First Bridging Workshop, 2007 Nov. 11–15, Aleppo, Syria. International Center for Agricultural Research in the Dry Areas. Available from: http://www.ais.unwater.org/ais/pluginfile.php/225/mod_label/intro/2008%20Proc%20First%20Bridging%20Workshop%20%283%29.pdf. Accessed 2014 August 14.
- Rai PK, Tripathi BD. 2007. Microbial contamination in vegetables due to irrigation with partially treated municipal wastewater in a tropical city. *Intl J Environ Health Res* 17:389–95.
- Raschid-Sally L, Jayakody P. 2009. Drivers and characteristics of wastewater agriculture in developing countries: results from a global assessment. Available from: http://www.iwmi.cgiar.org/Publications/IWMI_Research_Reports/PDF/PUB127/RR127.pdf. Accessed 2014 August 14.
- Rechenburg A, Koch C, Classen T, Kistemann T. 2006. Impact of sewage treatment plants and combined sewer overflow basins on the microbiological quality of surface water. *Water Sci Technol* 54:95–9.
- Reid DC, Edwards AC, Cooper D, Wilson E, McGaw BA. 2003. The quality of drinking water from private water supplies in Aberdeenshire, UK. *Water Res* 37:245–54.
- Reingold, AL. 1998. Outbreak investigations—a perspective. *Emerg Infect Dis* 4:21. Available from: http://wwwnc.cdc.gov/eid/article/4/1/98-0104_article. Accessed 2014 August 14.
- Rhodes MW, Kator HOWA. 1988. Survival of *Escherichia coli* and *Salmonella* spp. in estuarine environments. *Appl Environ Microbiol* 54:2902–7.
- Rosenkötter N, Ziemann A, Krafft T, Riesgo LGC, Vergeiner G, Brand H. 2014. Noninfectious events under the International Health Regulations (2005) in Europe: a case for syndromic surveillance. *J Public Health Policy* 35:311–26.
- Savichtcheva O, Okabe S. 2006. Alternative indicators of fecal pollution: relations with pathogens and conventional indicators, current methodologies for direct pathogen monitoring and future application perspectives. *Water Res* 40:2463–76.
- Szakli E, Alexopoulos A, Leontinidis M. 2007. Rainwater harvesting, quality assessment and utilization in Kefalonia Island, Greece. *Water Res* 41:2039–47.
- Scallan E, Griffin PM, Angulo FJ, Tauxe RV, Hoekstra RM. 2011. Foodborne illness acquired in the United States—unspecified agents. *Emerging Infectious Diseases* 17. Available from: http://wwwnc.cdc.gov/eid/article/17/1/p2-1101_article. Accessed 2014 August 14.
- SCF. 2002. Risk profile on the microbiological contamination of fruits and vegetables eaten raw. Report of the scientific committee on food. Available from: http://ec.europa.eu/food/fs/sc/scf/out125_en.pdf. Accessed 2014 August 14.
- Schets, FM, During M, Italiaander R, Heijnen L, Rutjes SA, Van der Zwaluw WK, de Roda Husman AM. 2005. *Escherichia coli* O157:H7 in drinking water from private water supplies in the Netherlands. *Water Res* 39:4485–93.
- Scott C, Drechsel P, Raschid-Sally L, Bahri A, Mara D, Redwood M, Jiménez B. 2010. Wastewater irrigation and health: challenges and outlook for mitigating risks in low-income countries. In: Drechsel P, Scott C, Raschid-Sally L, Redwood M, Bahri A, editors. *Wastewater irrigation and health: assessing and mitigating risk in low-income countries*. London: Earthscan. p 381–94.
- Seo KH, Frank JF. 1999. Attachment of *Escherichia coli* O157:H7 to lettuce leaf surface and bacterial viability in response to chlorine treatment as demonstrated by using confocal scanning laser microscopy. *J Food Prot* 62:3–9.
- Shah K, Sharma PK, Nandi I, Singh N. 2014. Water sustainability: reforming water management in new global era of climate change. *Environ Sci Pollut Res* 21:11603–4.
- Sivapalasingam S, Friedman CR, Cohen L, Tauxe RV. 2004. Fresh produce: a growing cause of outbreaks of foodborne illness in the United States, 1973 through 1997. *J Food Prot* 67:2342–53.
- Sliva L, Dudley Williams D. 2001. Buffer zone versus whole catchment approaches to studying land use impact on river water quality. *Water Res* 35:3462–72.
- Söderström A, Österberg P, Lindqvist A, Jönsson B, Lindberg A, Blide Ulander S, Welinder-Olsson C, Löfdahl S, Kaijser B, De Jong B. 2008. A large *Escherichia coli* O157 outbreak in Sweden associated with locally produced lettuce. *Foodborne Pathog Dis* 5:339–49.
- Solomon EB, Pang HJ, Matthews KR. 2003. Persistence of *Escherichia coli* O157:H7 on lettuce plants following spray irrigation with contaminated water. *J Food Protect* 66:2198–202.
- Solomon EB, Yaron S, Matthews KR. 2002. Transmission of *Escherichia coli* O157:H7 from contaminated manure and irrigation water to lettuce plant tissue and its subsequent internalization. *Appl Environ Microbiol* 68:397–400.
- Song I, Stine SW, Choi CY, Gerba CP. 2006. Comparison of crop contamination by microorganisms during subsurface drip and furrow irrigation. *J Environ Eng* 132:1243–8.
- SQF Institute. 2010. SQF 1000 Code: A HACCP based supplier assurance code for the primary producer. Available from: <http://www.sqfi.com/wp-content/uploads/SQF-1000-Code.pdf>. Accessed 2014 August 14.
- Steele M, Odumeru J. 2004. Irrigation water as source of foodborne pathogens on fruit and vegetables. *J Food Prot* 67:2839–49.
- Stine SW, Song I, Choi CY, Gerba CP. 2005. Application of microbial risk assessment to the development of standards for enteric pathogens in water used to irrigate fresh produce. *J Food Prot* 68:913–8.
- Strawn LK, Schneider KR, Danyluk MD. 2011. Microbial safety of tropical fruits. *Crit Rev Food Sci Nutr* 51:132–45.
- Stumpf CH, Piehler MF, Thompson S, Noble RT. 2010. Loading of fecal indicator bacteria in North Carolina tidal creek headwaters: hydrographic patterns and terrestrial runoff relationships. *Water Res* 44:4704–15.
- Sugumar G, Mariappan S. 2003. Survival of *Salmonella* sp. in freshwater and seawater microcosms under starvation. *Asian Fisheries Sci* 16:247–56.
- Suslow TV, Oriá MP, Beuchat LR, Garrett EH, Parish ME, Harris LJ, Farber JN, Busta FF. 2003. Production practices as risk factors in microbial food safety of fresh and fresh cut produce. *Compr Rev Food Sci Food Saf* 2:38–77.
- Szewzyk U, Szewzyk R, Manz W, Schleifer KH. 2000. Microbiological safety of drinking water. *Annu Rev Microbiol* 54:81–127.
- Theofel CG, Harris LJ. 2009. Impact of preinoculation culture conditions on the behavior of *Escherichia coli* O157:H7 inoculated onto romaine lettuce (*Lactuca sativa*) plants and cut leaf surfaces. *J Food Prot* 72:1553–9.
- Tyrrel SF, Knox JW, Weatherhead EK. 2006. Microbiological water quality requirements for salad irrigation in the United Kingdom. *J Food Prot* 69:2029–35.
- United Fresh Produce Association. 2002. Packaging design for fresh-cut produce. Available from: http://www2.unitedfresh.org/forms/store/ProductFormPublic/search?action=1&Product_categories_Checkboxes_exists=1&Product_categories_Checkboxes=11 accessed 2014 August 14.
- United Fresh Produce Association. 2005. Commodity specific food safety guidelines for the melon supply chain. Available from: http://www2.unitedfresh.org/forms/store/ProductFormPublic/search?action=1&Product_categories_Checkboxes_exists=1&Product_categories_Checkboxes=11

- United Fresh Produce Association. 2008. Commodity specific food safety guidelines for the fresh tomatoes supply chain. Available from: http://www2.unitedfresh.org/forms/store/ProductFormPublic/search?action=1&Product_categories_Checkboxes_exists=1&Product_categories_Checkboxes=11
- United Fresh Produce Association. 2010. Best practices for garment use in fresh-cut produce plants. Available from: http://www2.unitedfresh.org/forms/store/ProductFormPublic/search?action=1&Product_categories_Checkboxes_exists=1&Product_categories_Checkboxes=11
- US Environmental Protection Agency. 2003. Bacterial water quality standards for recreational waters (freshwater and marine waters). Available from: http://water.epa.gov/type/oceb/beaches/upload/2003_06_19_beaches_local_statrept.pdf. Accessed 2014 August 14.
- US Environmental Protection Agency. 2004. Guidelines for water reuse. Available from: <http://water.epa.gov/aboutow/own/upload/Water-Reuse-Guidelines-625r04108.pdf>. Accessed 2014 August 14.
- Van Haute S, Sampers I, Jacxsens L, Uyttendaele M. 2013. Selection criteria for water disinfection techniques in agricultural practices. *Critical Reviews in Food Science and Nutrition* (epublished). Available from: <http://www.tandfonline.com/doi/pdf/10.1080/10408398.2012.70536>. Accessed 24 August 14. doi: 10.1080/10408398.2012.705360.
- Verbeten E. 1998. Irrigation in arid and semi-arid environments. IMAROM Working Paper Series No. 1. Available from: [http://www.heindehaas.com/IMAROM/IMAROM%20working%20papers/IMAROM%20Working%20Paper%2001%20\(Verbeten\).pdf](http://www.heindehaas.com/IMAROM/IMAROM%20working%20papers/IMAROM%20Working%20Paper%2001%20(Verbeten).pdf). Accessed 2014 August 14.
- VMM. 2013. Heffingen op grondwaterverbruik. Available from: "<https://www.vmm.be/water/heffingen%20accessed%20on%2011/12/2013>" <https://www.vmm.be/water/heffingen>. Accessed 2013 December 11.
- Wang G, Doyle MP. 1998. Survival of enterohemorrhagic *Escherichia coli* O157:H7 in water. *J Food Prot* 61:662–7.
- Warriner K, Namvar A. 2010. The tricks learnt by human enteric pathogens from phytopathogens to persist within the plant environment. *Curr Opin Biotechnol* 21:131–6.
- Wood JD, Bezanson GS, Gordon RJ, Jamieson R. 2010. Population dynamics of *Escherichia coli* inoculated by irrigation into the phyllosphere of spinach grown under commercial production conditions. *Intl J Food Microbiol* 143:198–204.
- World Health Organization. 1989. Health guidelines for the use of wastewater in agriculture and aquaculture. Available from: http://whqlibdoc.who.int/trs/WHO_TRS_778.pdf. Accessed 2014 August 14.
- World Health Organization. 2006. WHO guidelines for the safe use of wastewater, excreta and grey water. Wastewater use in agriculture. Available from: http://whqlibdoc.who.int/publications/2006/9241546832_eng.pdf. Accessed 2014 August 14.
- World Health Organization. 2008a. Annex 2: Decision instrument for the assessment and notification of events that may constitute a public health emergency of international concern in international health regulations. *International Health Regulations (2005)*. World Health Organization. p 43–6.
- World Health Organization. 2008b. Microbiological risk assessment series. Microbiological hazards in fresh fruits and vegetables. Available from: http://www.who.int/foodsafety/publications/micro/MRA_FruitVegetables.pdf. Accessed 2014 August 14.
- World Health Organization. 2010. Guidelines for drinking-water quality. Available from: http://www.who.int/water_sanitation_health/dwq/guidelines/en/. Accessed 2014 August 14.
- World Health Organization and UNICEF. 2000. Global water supply and sanitation assessment 2000 report. Available from: http://www.who.int/water_sanitation_health/monitoring/jmp2000.pdf. Accessed 2014 August 14.
- Yaziz MI, Gunting H, Sapari N, Ghazali AW. 1989. Variations in rainwater quality from roof catchments. *Water Res* 23:761–5.
- Yengoh GT, Tchuente A, Armah FA, Odoi JO. 2010. Impact of prolonged rainy seasons on food crop production in Cameroon. *Mitig Adapt Strateg Glob Change* 15:825–41.
- Yermiyahu U, Tal A, Ben-Gal A, Bar-Tal A, Tarchisky J, Lahav O. 2007. Environmental science. Rethinking desalinated water quality and agriculture. *Science* 318:920–1.
- Zhang G, Ma L, Phelan VH, Doyle MP. 2009. Efficacy of antimicrobial agents in lettuce leaf processing water for control of *Escherichia coli* O157:H7. *J Food Prot* 72:1392–7.